

Multi-messenger light curves from gamma-ray bursts

Mauricio Bustamante


Center for Cosmology and AstroParticle Physics (CCAPP)
The Ohio State University

NSSTC Space Seminar
Huntsville — August 02, 2016




Ultra-high-energy cosmic rays

After 50+ years of UHECR measurements —



GAISSER, STANEV, TILAV,
Front. Phys. China 8, 748 (2013)

The electromagnetic sky




B. LACKI,
MNRAS 406, 863 (2010)

A new player: high-energy astrophysical neutrinos


The era of neutrino astronomy has begun!

IceCube has reported 54 events with 30 TeV – 2 PeV in 4 years



The newest kid on the block: gravitational waves


We have started listening to compact object mergers —



LIGO & VIRGO, *PRL* 116, 061102 (2016)


What about the sources of HE neutrinos?

Arrival directions compatible with an **isotropic** distribution –



No association with sources found **yet**

The multi-messenger picture



We will use gamma-ray bursts (GRBs) as a concrete source example

Why are GRBs interesting as potential ν sources?

- 1 They are the best candidates for detection of coincident high-energy e.m.–neutrino emission at PeV ν energies
- 2 Neutrinos from GRB afterglows are expected to be important at EeV energies (observable by radio neutrino detectors)

[K. MURASE, *PRD* **76**, 123001 (2007)] [S. RAZZAQUE, L. YANG, *PRD* **91**, 043003 (2015)]

- 3 Dark, “failed” GRBs might contribute an important part to the diffuse flux seen by IceCube


[P. MÉSZAROS, E. WAXMAN, *PRL* **87**, 171102 (2001)]

Here we will focus on issue 1
and explore the prompt emission of neutrinos in GRBs

Why is *now* a good time to do this?

better, bigger detectors + loads of data + bright future

Neutrinos



- ▶ IceCube: diffuse flux of HE astrophysical ν 's
- ▶ No point sources yet
- ▶ GRBs: low bg due to time and direction cuts
- ▶ IceCube-Gen2

GRBs



- ▶ *Fermi*: ~ 250 GRBs yr^{-1} in 8 keV – 40 MeV
- ▶ ~ 12 GRBs yr^{-1} in 20 MeV – 300 GeV
- ▶ different wavelengths: INTEGRAL, *Swift*
- ▶ 1000's GRBs detected so far

UHECRs



- ▶ Auger: 69 events > 57 EeV
- ▶ Telescope Array: 72 events
- ▶ surface + fluorescence
- ▶ future: LHAASO, JEM-EUSO

GRBs – good candidates for UHE CR & ν sources

GRBs are among the best candidate sources for CRs *and* ν 's:

- ▶ radiated energy of $\sim 10^{52} - 10^{53}$ erg
- ▶ intense magnetic fields of $\sim 10^5$ G
- ▶ magnetically-confined p 's shock-accelerated to $\sim 10^{12}$ GeV
- ▶ plus: low backgrounds (for ν 's) due to small time window

Problem: experiments (IceCube, ANTARES) are starting to strongly constrain the simplest joint emission models

Solution: we need to build more realistic models!

GRBs – good candidates for UHE CR & ν sources

GRBs are among the best candidate sources

- ▶ radiated energy of $\sim 10^{52} - 10^{53}$ erg
- ▶ intense magnetic fields of $\sim 10^5$ G
- ▶ magnetically-confined p 's shock-accelerated to $\sim 10^{12}$ GeV
- ▶ plus: low backgrounds (for ν 's) due to small time window

10^{20} erg	H bomb
10^{26} erg	killer asteroid
10^{40} erg	Death Star
10^{33} erg s $^{-1}$	Sun
10^{41} erg s $^{-1}$	supernova
10^{45} erg s $^{-1}$	galaxy


Problem: experiments (IceCube, ANTARES) are starting to strongly constrain the simplest joint emission models

Solution: we need to build more realistic models!

GRBs – what are they?


GRBs: the most luminous explosions in the Universe

- ▶ **brief** flashes of gamma rays:
from 0.1 s to few 100's s
- ▶ isotropically distributed in the sky
- ▶ they are **far**: most occur
at ~ 1 Gpc from us ($z \approx 2$)
- ▶ they are **rare**: $\sim 0.3 \text{ Gpc}^{-3} \text{ yr}^{-1}$
- ▶ two populations:
 - ▶ **short-duration** (< 2 s): neutron star–neutron star or NS–black hole mergers
 - ▶ **long-duration** (> 2 s): associated to hypernovae
- ▶ powered by matter accretion
onto a black hole



GRBs – a zoo of light curves


GRB light curves come in different shapes:



variability timescale (width of pulses) $\equiv t_v \approx 1$ ms


GRBs explained – the fireball model

Developed by Mészáros, Reese, Goodman, Pachinsky, *et al.* in the 1990s





Internal collisions

Relativistically-expanding blobs of plasma collide with each other, merge, and emit UHE particles —



Producing the UHE ν 's, CRs, γ rays – a first look

Joint production of UHECRs, ν 's, and γ 's:



After propagation, with flavor mixing:

$$\nu_e : \nu_\mu : \nu_\tau : p = 1 : 1 : 1 : 1$$

("one ν_μ per cosmic ray")

This **neutron model** of CR emission is now strongly disfavoured

IceCube, *Nature* **484**, 351 (2012)

M. AHLERS *et al.* *Astropart. Phys.* **35**, 87 (2011)

What are the ingredients?

To calculate the ν flux from a GRB, we need:

- ▶ its gamma-ray luminosity L_{γ}^{iso} [erg s $^{-1}$] **[measured]**
- ▶ its variability timescale t_{ν} [s], from the light curve **[measured]**
- ▶ the break energy of its photon spectrum $\epsilon_{\gamma, \text{break}}$ [MeV] **[measured]**
- ▶ the bulk Lorentz factor of its jet Γ **[estimated]**
- ▶ the energy in electrons, magnetic field, protons **[estimated]**

Now let us cook up the neutrinos ►

Normalizing neutrinos with observed gamma rays

$$\text{energy in neutrinos} \propto \text{energy in gamma rays}$$

In detail, for each GRB,

$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \underbrace{\left[1 - (1 - \langle x_{p \rightarrow \pi} \rangle)^{\Delta R / \lambda_{p\gamma}} \right]}_{f_\pi} \frac{1}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} d\epsilon_\gamma \epsilon_\gamma F_\gamma(\epsilon_\gamma)$$

f_π : fraction of total proton energy transferred to pions

ΔR : size of the emitting region

$\lambda_{p\gamma}$: mean free path for $p\gamma$ interactions

$\langle x_{p \rightarrow \pi} \rangle$: avg. fraction of p energy transferred to a π in one interaction

f_e^{-1} : ratio of energy in protons to energy in photons/electrons

$$\frac{\Delta R}{\lambda_{p\gamma}} = \left(\frac{L_\gamma^{\text{iso}}}{10^{52} \text{ erg s}^{-1}} \right) \left(\frac{0.01}{t_\nu} \right) \left(\frac{10^{2.5}}{\Gamma} \right)^4 \left(\frac{\text{MeV}}{\varepsilon_{\gamma, \text{break}}} \right)$$

The original recipe: conventional fireball model

Observed gamma-ray fluence [$\text{GeV}^{-1} \text{ cm}^{-2}$]

$$\mathcal{F}_\gamma(\varepsilon_\gamma) \propto \begin{cases} (\varepsilon_\gamma/\varepsilon_{\gamma,\text{break}})^{-1} & , \varepsilon_\gamma < \varepsilon_{\gamma,\text{break}} = 1 \text{ MeV} \\ (\varepsilon_\gamma/\varepsilon_{\gamma,\text{break}})^{-2.2} & , \varepsilon_\gamma \geq \varepsilon_{\gamma,\text{break}} \end{cases}$$

+

Assumed proton spectrum in the source


$$N'_p(E_p) \propto E_p'^{-2}$$

=

Neutrino spectrum from $p\gamma$, via Δ resonance

$$F_\nu(E_\nu) \propto \begin{cases} \left(\frac{E_\nu}{E_{\nu,\text{break}}}\right)^{-\alpha_\nu} & , E_\nu < E_{\nu,\text{break}} \\ \left(\frac{E_\nu}{E_{\nu,\text{break}}}\right)^{-\beta_\nu} & , E_{\nu,\text{break}} \leq E_\nu < E_{\nu,\mu} \\ \left(\frac{E_\nu}{E_{\nu,\text{break}}}\right)^{-\beta_\nu} \left(\frac{E_\nu}{E_{\nu,\mu}}\right)^{-2} & , E_\nu \geq E_{\nu,\mu} \end{cases}$$

Neutrino spectrum – conventional fireball




E. WAXMAN, J. N. BAHCALL, *PRL* **78**, 2292 (1997)

D. GUETTA *et al.*, *Astropart. Phys.* **20**, 429 (2004)

The *one-collision* burst assumption

All internal collisions are identical and occur at the **same** radius —




- ▶ Average speed Γ inferred from afterglow observations
- ▶ “Variability timescale” t_v measured from the light curve
- ▶ Redshift z measured for the host galaxy
- ▶ “Isotropically-equivalent volume”: $V'_{\text{iso}} = 4\pi R_C^2 l'$
- ▶ $N_{\text{coll}} \approx T_{90} / t_v \sim 100\text{--}1000$ identical collisions

Neutron model of UHECR emission under tension?

In 2012, IceCube ruled this analytical version of the fireball model –

- ▶ assumed a fixed baryonic loading of 10
- ▶ extrapolated diffuse ν flux from 117–215 GRBs (“quasi-diffuse”)
- ▶ **analytical calculation** – in tension with upper bounds



IceCube, *Nature* 484, 351 (2012)

M. AHLERS *et al.* *Astropart. Phys.* 35, 87 (2011)

D. GUETTA *et al.* *Astropart. Phys.* 20, 429 (2004)

NeuCosmA: (revised) GRB particle emission – I

Two ingredients:

$$\underbrace{N'_p(E'_p)}_{\text{proton density at the source } [\text{GeV}^{-1} \text{ cm}^{-3}]} \underset{\text{NeuCosmA}}{\otimes} \underbrace{N'_\gamma(E'_\gamma)}_{\text{photon density at the source}} = \underbrace{Q'_\nu(E'_\nu)}_{\text{emitted neutrino spectrum } [\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}]}$$

► Photons (same shape as observed at Earth):

$$N'_\gamma(E'_\gamma) = \begin{cases} (E'_\gamma/E'_{\gamma,\text{break}})^{-1} & , E'_{\gamma,\text{min}} = 0.2 \text{ eV} \leq E'_\gamma < E'_{\gamma,\text{break}} = 1 \text{ keV} \\ (E'_\gamma/E'_{\gamma,\text{break}})^{-2.2} & , E'_\gamma \geq E'_{\gamma,\text{break}} \\ 0 & , \text{otherwise} \end{cases}$$

► Protons: $N'_p(E'_p) \propto E_p'^{-\alpha_p} e^{-E'_p/E'_{p,\text{max}}} \quad (\alpha_p \gtrsim 2)$

NeuCosmA: (revised) GRB particle emission – I

Two ingredients:

$$\underbrace{N'_p(E'_p)}_{\text{proton density at the source } [\text{GeV}^{-1} \text{ cm}^{-3}]} \underset{\text{NeuCosmA}}{\otimes} \underbrace{N'_\gamma(E'_\gamma)}_{\text{photon density at the source}} = \underbrace{Q'_\nu(E'_\nu)}_{\text{emitted neutrino spectrum } [\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}]}$$

► Photons (same shape as observed at Earth):

$$N'_\gamma(E'_\gamma) = \begin{cases} (E'_\gamma/E'_{\gamma,\text{break}})^{-1} & , E'_{\gamma,\text{min}} = 0.2 \text{ eV} \leq E'_\gamma < E'_{\gamma,\text{break}} = 1 \text{ keV} \\ (E'_\gamma/E'_{\gamma,\text{break}})^{-2.2} & , E'_\gamma \geq E'_{\gamma,\text{break}} \\ 0 & , \text{otherwise} \end{cases}$$

$$t'_{\text{acc}}(E'_{p,\text{max}}) = \min \left[t'_{\text{dyn}}, t'_{\text{syn}}(E'_{p,\text{max}}), t'_{p\gamma}(E'_{p,\text{max}}) \right]$$

► Protons: $N'_p(E'_p) \propto E_p'^{-\alpha_p} e^{-E'_p/E'_{p,\text{max}}} \quad (\alpha_p \gtrsim 2)$

NeuCosmA: (revised) GRB particle emission – II

Normalize the particle densities at the source —

► Photons:

$$\underbrace{\text{photon energy density per collision}}_{\int E'_\gamma N'_\gamma(E'_\gamma) dE'_\gamma} = \frac{\overbrace{E_{\gamma, \text{tot}}^{\text{iso}, \prime} \sim 10^{53} \text{ erg (from observed fluence)}}^{\text{total gamma-ray energy of burst}}}{\underbrace{N_{\text{coll}}}_{\text{number of collisions}} \cdot \underbrace{V'_{\text{iso}}}_{\text{volume of one collision}}}$$

► Protons: *baryonic loading* (energy in p 's / energy in e 's + γ 's), e.g., 10

$$\underbrace{\text{proton energy density per collision}}_{\int E'_p N'_p(E'_p) dE'_p} = \frac{1}{f_e} \cdot \text{photon energy density per collision}$$

NeuCosmA: (revised) GRB particle emission – III

Injected/ejected spectrum of secondaries (π , K , n , ν , etc.):

$$Q'(E') = \int_{E'}^{\infty} \frac{dE'_p}{E'_p} N'_p(E'_p) \int_0^{\infty} c \, dE'_\gamma \, N'_\gamma(E'_\gamma) \, R(x, y)$$


response function

$x \equiv E'/E'_p \quad y \equiv E'_p E'_\gamma / (m_p c^2)$


R contains cross sections, multiplicities for different channels

What does NeuCosmA include?

- ▶ $p\gamma \rightarrow \Delta^+ (1232) \rightarrow \pi^0, \pi^+, \dots$
- ▶ extra K , n , π^- , multi- π prod. modes
- ▶ synchrotron losses of secondaries
- ▶ adiabatic cooling
- ▶ full photon spectrum
- ▶ neutrino flavor transitions



NeuCosmA – how the neutrino spectrum changes



For example, GRB080603A:

1. Correction to analytical model (IC-FC → RFC)
2. Change due to full numerical calculation

IC-FC: IceCube-Fireball Calculation
RFC: Revised Fireball Calculation
NFC: Numerical Fireball Calculation


The new prediction of the quasi-diffuse GRB ν flux

Repeat the IceCube GRB neutrino analysis, with NeuCosmA —

- ▶ Same GRB sample and parameters
- ▶ Calculate ν fluence for each burst and stacked fluence $F_\nu(E_\nu)$
- ▶ Quasi-diffuse flux ($N_{\text{GRB}} = 117$):

$$\phi_\nu(E_\nu) = F_\nu(E_\nu) \frac{1}{4\pi} \frac{1}{N_{\text{GRB}}} \frac{667 \text{ bursts}}{\text{yr}}$$



Flux ~ 1 order of magnitude lower!



S. HÜMNER, P. BAERWALD, W. WINTER,
PRL 108, 231101 (2012)

Improved IceCube limits (2016)



- ▶ 3 yr of showers (all flavors) + 4 yr of upgoing tracks with > 1 TeV
- ▶ Larger GRB catalogue (807 bursts)
- ▶ Six coincident events found (five showers + one track)
- ▶ Low statistical significance
- ▶ $\lesssim 1\%$ of the diffuse flux can be from prompt GRB emission



ICECUBE, 1601.06484

Improved IceCube limits (2016)


- ▶ 3 yr of showers (all flavors) + 4 yr of upgoing tracks with > 1 TeV
- ▶ Larger GRB catalogue (807 bursts)
- ▶ Six coincident events found (five showers + one track)
- ▶ Low statistical significance
- ▶ $\lesssim 1\%$ of the diffuse flux can be from prompt GRB emission



ICECUBE, 1601.06484

What about low-luminosity and choked GRBs?

- ▶ Low-luminosity and choked GRBs might be in the same family as high-luminosity long GRBs
- ▶ Due to lower jet speeds (Γ_b), they do not break out
- ▶ They might explain the TeV region of the IceCube diffuse ν flux:



I. TAMBORRA, S. ANDO, PRD 93, 053010 (2016)

Going beyond the neutron model

The neutron model hinges on:

- ① p 's magnetically confined, only n 's escape
- ② p 's interact at most once, n 's do not (*optically thin source*)

However, under the “one ν_μ per CR” hypothesis, GRBs **are disfavored** as the sole sources of UHECRs ([AHLERS *et al.*](#)).

[M. AHLERS, M. GONZÁLEZ-GARCÍA, F. HALZEN, *Astropart. Phys.* **35**, 87 \(2011\)](#)

Going beyond the neutron model

The neutron model hinges on:

- ➊ p 's magnetically confined, only n 's escape
- ➋ p 's interact at most once, n 's do not (*optically thin source*)

However, under the “one ν_μ per CR” hypothesis, GRBs **are disfavored** as the sole sources of UHECRs ([AHLERS *et al.*](#)).

What if ➊ and ➋ are violated?

- ▶ p 's “leak out”, not accompanied by (direct) ν production
- ▶ multiple p interactions enhance the ν flux
- ▶ in *optically thick sources*, only n 's at the borders escape


[M. AHLERS, M. GONZÁLEZ-GARCÍA, F. HALZEN, *Astropart. Phys.* **35**, 87 \(2011\)](#)

Going beyond the neutron model

We have improved the model – now UHECRs escape as either:

- ▶ **neutrons**, which decay into protons outside the source; or
- ▶ **protons** that leak out without interacting inside the source


Relative contributions determined by $\tau_n \equiv \left(t_{p\gamma}^{-1} / t_{\text{dyn}}^{-1} \right) \Big|_{E'_{p,\text{max}}}$




A two-component model of UHECR emission

Sample neutrino fluences –

Optically **thin** source




Optically **thick** source




P. BAERWALD, MB, W. WINTER, *ApJ* 768, 186 (2013)


From the sources to us



From the sources to us




From the sources to us



From the sources to us


Because of the cosmological expansion:

$$\text{energy at Earth} = \frac{\text{energy at production time}}{1+z}$$




From the sources to us


Cosmological photon backgrounds:



◀ they evolve with redshift,
i.e., $n_\gamma(\epsilon, z)$



From the sources to us




From the sources to us

γ 's and e^\pm 's dump energy into e.m. cascades through

- ▶ pair production, $\gamma + \gamma_b \rightarrow e^+ + e^-$
- ▶ inverse Compton scattering, $e^\pm + \gamma_b \rightarrow e^\pm + \gamma$

Lower-energy (GeV–TeV) gamma-rays detected by Fermi-LAT



From the sources to us

p 's are deflected by extragalactic magnetic fields


⇒ except for the most energetic ones, they are
not expected to point back to the sources

Pierre Auger found weak correlation
with known AGN positions


They lose energy through:

- ▶ pair production, $p + \gamma_b \rightarrow p + e^+ + e^-$
- ▶ photohadronic interactions, $p\gamma_b$

depend on the redshift evolution
of the cosmological γ backgrounds



From the sources to us



Initial UHE ν flavor fluxes: $\nu_e : \nu_\mu : \nu_\tau = 1 : 2 : 0$

Probability of $\nu_\alpha \rightarrow \nu_\beta$ transition: $P_{\alpha\beta}(E_0, z)$

Flavor oscillations redistribute the fluxes


– at Earth: $\nu_e : \nu_\mu : \nu_\tau \approx 1 : 1 : 1$ (might be changed by exotic physics!)

MB, Beacom, Winter, *PRL* 115, 161302 (2015)


What do the diffuse fluxes look at Earth, then?

Neutron model vs. two-component model:
prompt and cosmogenic ν 's

UHECRs



Neutrinos




P. BAERWALD, MB, W. WINTER, *ApJ* 768, 186 (2013)

P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* 62, 66 (2015)

See also: H. HE *et al.*, *ApJ* 752, 29 (2012)

Constraints from experimental data

Fit the UHECR flux to Telescope Array data & enforce the IceCube GRB ν and cosmogenic ν upper limits —



P. BAERWALD, MB, AND W. WINTER,
Astropart. Phys. **62**, 66 (2015)

More relativistic (higher Γ) GRBs are needed

Going further: an evolving burst


Consider an evolving fireball –

- ▶ the fireball expands with time
- ▶ ~ 1000 shells propagate with different speeds
- ▶ they have different masses
- ▶ they collide at different radii
(collisions no longer identical)

What is different?

The particle (γ , p) densities fall as the fireball expands – particle production conditions change with time/radius

S. KOBAYASHI, T. PIRAN, R. SARI, *ApJ* 490, 92 (1997)
F. DAIGNE, R. MOCHKOVITCH, *MNRAS* 296, 275 (1998)




Initial distribution of shell speeds

Distribution of initial shell speeds (Lorentz factors):

$$\ln \left(\frac{\Gamma_{k,0} - 1}{\Gamma_0 - 1} \right) = A_\Gamma \cdot x$$

x follows a Gaussian distribution, $P(x) dx = dx e^{-x^2/2} / \sqrt{2\pi}$



$$A_\Gamma < 1$$

Speeds too similar, collisions only at large radii

$$A_\Gamma \gg 1$$


Spread too large, too many collisions at low radii

$$A_\Gamma \approx 1$$

Just right, burst has high efficiency of conversion of kinetic to radiated energy


Anatomy of an internal collision

1 Propagation




Anatomy of an internal collision

1 Propagation




2 Collision




Anatomy of an internal collision

1 Propagation




2 Collision



$$m_m = \frac{I_f m_f + I_s m_s}{I_m}$$

$$I_m < I_f, I_s$$

3 Radiation



Part of the initial kinetic energy radiated as γ 's, ν 's, p 's, and n 's:

$$E_{\text{coll}}^{\text{iso}} = (E_{\text{kin},f}^{\text{iso}} - E_{\text{kin},m}^{\text{iso}}) + (E_{\text{kin},m}^{\text{iso}} - E_{\text{kin},s}^{\text{iso}})$$

$$\underbrace{\epsilon_e}_{1/12} E_{\text{coll}}^{\text{iso}}$$

energy in photons

$$\underbrace{\epsilon_B}_{1/12} E_{\text{coll}}^{\text{iso}}$$

energy in magnetic fields

$$\underbrace{\epsilon_p}_{5/6} E_{\text{coll}}^{\text{iso}}$$

energy in baryons


Tracking each collision individually

Each collision occurs in a different emission regime –

Sub-photospheric: $\tau_{e\gamma} > 1$

$\nu_\mu + \bar{\nu}_\mu$ fluence


neutrinos



(observer's frame)

maximum p energy

cosmic rays




(source frame)

Limited by $\gamma + \gamma \rightarrow e^+ + e^-$



maximum γ energy

gamma-rays



Synthetic light curves


An emission pulse is assigned to each collision
– their superposition yields a **synthetic light curve**:



MB, P. BAERWALD, K. MURASE, W. WINTER
Nature Commun. **6**, 6783 (2015)


Synthetic light curves

An emission pulse is assigned to each collision
– their superposition yields a **synthetic light curve**:



Different particles come from different jet regions

Emission of different species peaks at different collision radii –



Why?

As the fireball expands, photon and proton densities fall


Why does it matter?

GRB parameters derived from gamma-ray observations might not be adequate to describe ν and UHECR emission

MB, P. BAERWALD, K. MURASE, W. WINTER, *Nat. Commun.* **6**, 6783 (2015)

A robust minimal diffuse ν flux from GRBs

- ▶ Take the simulated burst as stereotypical
- ▶ Quasi-diffuse neutrino flux, assuming 667 identical GRBs per year:




MB, P. BAERWALD, K. MURASE, W. WINTER
Nature Commun. **6**, 6783 (2015)

What makes a GRB bright in neutrinos?

- ▶ **Goal:** to use the morphology of the GRB gamma-ray light curve to assess whether the burst is neutrino-bright
- ▶ The central engine determines the features of the light curve


Undisciplined GRB engine

- ▶ Broad Γ distribution
- ▶ *E.g.*, engine emits shells with log-normal Γ distrib.



Disciplined GRB engine

- ▶ Narrow Γ distribution
- ▶ *E.g.*, engine emits shells with oscillating Γ





Light curves

Undisciplined GRB engine

- ▶ Fast variability dominates
- ▶ No broad pulses

Disciplined GRB engine

- ▶ Broad pulses dominate
- ▶ Fast variability on top





How many optically thick collisions?

Undisciplined GRB engine

- ▶ Shells have very different speeds
- ▶ Collide quickly, close to engine
- ▶ High ρ and γ densities
- ▶ ~ 10 optically-thick bursts near the photosphere

Disciplined GRB engine

- ▶ Shells have similar speeds
- ▶ Collide far from engine
- ▶ Low ρ and γ densities
- ▶ All (superphotospheric) collisions are optically thin




So which burst is neutrino-bright?

Compare the quasi-diffuse neutrino fluxes:


Undisciplined GRB engine

$$\sim 10^{-11} \text{ GeV cm}^{-1} \text{ s}^{-1} \text{ sr}^{-1}$$



Disciplined GRB engine


$$\sim 5 \cdot 10^{-13} \text{ GeV cm}^{-1} \text{ s}^{-1} \text{ sr}^{-1}$$



∴ An undisciplined engine makes a GRB neutrino-bright

Time delays in gamma-ray light curves

Neutrino-weak bursts show time delays in different energy bands —



MB, MURASE, WINTER, 1606.02325

Conclusions ... and the future


- ▶ GRBs *are* good UHECR and ν source candidates
- ▶ *But* the CR- ν - γ connection is trickier than originally thought
- ▶ Contribute to the diffuse HE astrophysical ν flux at the few % level
- ▶ Likely to be the first point neutrino sources to be resolved
- ▶ Need *next-gen* neutrino telescopes (IceCube-Gen2, KM3NeT) ...
- ▶ ... while Auger, Telescope Array, CTA, *etc.* gather extensive UHECR statistics
- ▶ Based on gamma-ray light curve morphology, we may build better GRB catalogues for ν searches

Backup slides

GRBs – an accidental discovery

After the 1963 Nuclear Test Ban Treaty, the U.S. launched six pairs of *Vela* satellites:


- ▶ They carried X-ray, gamma-ray, and neutron detectors
- ▶ *Vela* 5a-b had enough spatial resolution to pinpoint the direction of events
- ▶ Intense gamma-ray emission from a nuclear explosion lasts $\lesssim 10^{-6}$ s ...
- ▶ ... however, longer-lasting emissions were detected



VELA 5A/B SATELLITES (NASA)

GRBs – the first one detected


First GRB detected: July 2, 1967, 14:19 UTC



Detected by *Vela* 3, 4a, 4b (found on archival data)

What do they *look* like?


e.g., GRB060218 seen by *Swift*



SDSS, SWIFT COLLAB., SLOAN FOUNDATION, NSF, NASA

GRBs – two populations



Two populations of GRBs:



T_{90} : time during which 90% of gamma-ray energy is recorded

Improved IceCube bounds (2014)

- ▶ Only upgoing ν_μ 's with > 1 TeV used
- ▶ Four years of data (IC-40, -59, -79, -86)
- ▶ Larger GRB catalogue (506 bursts)
- ▶ One coincident event found, with low statistical significance
- ▶ $\lesssim 1\%$ of the diffuse flux can be from prompt GRB emission



[ICECUBE, *ApJ* 805, L5 (2015)]

A two-component model of CR emission

Optical depth:

$$\tau_n = \left. \frac{t_{p\gamma}^{-1}}{t_{\text{dyn}}^{-1}} \right|_{E_{p,\text{max}}} = \begin{cases} \lesssim 1, & \text{optically \textbf{thin} source} \\ > 1, & \text{optically \textbf{thick} source} \end{cases}$$

Particles can escape from within a shell of thickness λ'_{mfp} :

$$\left. \begin{array}{l} \lambda'_{p,\text{mfp}}(E') = \min [\Delta r', R'_L(E'), ct'_{p\gamma}(E')] \\ \lambda'_{n,\text{mfp}}(E') = \min [\Delta r', ct'_{p\gamma}(E')] \end{array} \right\} f_{\text{esc}} = \frac{\lambda'_{\text{mfp}}}{\Delta r'}$$


fraction of escaping particles

We *need* direct proton escape

Scan of the GRB emission parameter space –

acceleration efficiency $\longrightarrow \eta = 0.1$

$\eta = 1.0$




P. BAERWALD, MB, AND W. WINTER, *ApJ* 768, 186 (2013)

we need high efficiencies \Rightarrow direct proton escape *is* required


Star-formation-rate vs. GRB redshift evolution

The exclusion from cosmogenic ν 's grows if the number of GRBs evolves more strongly with redshift:

direct p escape, $\eta = 1.0$



direct p escape, $\eta = 1.0$




$$n_{\text{GRB}}(z) \propto \rho_{\text{SFR}}(z)$$

(star formation rate)

$$n_{\text{GRB}}(z) \propto \rho_{\text{SFR}}(z) \times (1+z)^{1.2}$$

NeuCosmA – the full photohadronic cross section




NeuCosmA – the full photohadronic cross section

Contributions to $(\nu_\mu + \bar{\nu}_\mu)$ flux from π^\pm decay divided in:

- ▶ $\Delta(1232)$ -resonance

P. BAERWALD, S. HÜMMER, AND W. WINTER,
Phys. Rev. D **83**, 067303 (2011)




NeuCosmA – the full photohadronic cross section

Contributions to $(\nu_\mu + \bar{\nu}_\mu)$ flux from π^\pm decay divided in:

- ▶ $\Delta(1232)$ -resonance
- ▶ Higher resonances

P. BAERWALD, S. HÜMMER, AND W. WINTER,
Phys. Rev. D83, 067303 (2011)




NeuCosmA – the full photohadronic cross section

Contributions to $(\nu_\mu + \bar{\nu}_\mu)$ flux from π^\pm decay divided in:

- ▶ $\Delta(1232)$ -resonance
- ▶ Higher resonances
- ▶ t -channel
(direct production)


P. BAERWALD, S. HÜMMER, AND W. WINTER,
Phys. Rev. D83, 067303 (2011)



NeuCosmA – the full photohadronic cross section

Contributions to $(\nu_\mu + \bar{\nu}_\mu)$ flux from π^\pm decay divided in:

- ▶ $\Delta(1232)$ -resonance
- ▶ Higher resonances
- ▶ t -channel
(direct production)
- ▶ High energy processes
(multiple π)



P. BAERWALD, S. HÜMMER, AND W. WINTER,
Phys. Rev. D **83**, 067303 (2011)

Especially "Multi π " contribution leads to **change of flux shape**; neutrino flux higher by up to a factor of 3 compared to WB treatment

NeuCosmA – further particle decays

$$\begin{aligned}\pi^+ &\rightarrow \mu^+ + \nu_\mu \\ \mu^+ &\rightarrow e^+ + \bar{\nu}_e + \bar{\nu}_\mu\end{aligned}$$

$$\begin{aligned}\pi^- &\rightarrow \mu^- + \bar{\nu}_\mu \\ \mu^- &\rightarrow e^- + \bar{\nu}_e + \nu_\mu\end{aligned}$$

$$K^+ \rightarrow \mu^+ + \nu_\mu$$

$$n \rightarrow p + e^- + \bar{\nu}_e$$

NeuCosmA – further particle decays


$$\pi^+ \rightarrow \mu^+ + \nu_\mu$$
$$\mu^+ \rightarrow e^+ + \bar{\nu}_e + \bar{\nu}_\mu$$

$$\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$$
$$\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$$

$$K^+ \rightarrow \mu^+ + \nu_\mu$$

$$n \rightarrow p + e^- + \bar{\nu}_e$$

Resulting ν_e flux (at the observer)



P. BAERWALD, S. HÜMMER, AND W. WINTER, *Phys. Rev. D* **83**, 067303 (2011)

NeuCosmA – further particle decays


$$\pi^+ \rightarrow \mu^+ + \nu_\mu$$
$$\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$$

$$\pi^- \rightarrow \mu^- + \bar{\nu}_\mu,$$
$$\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$$

$$K^+ \rightarrow \mu^+ + \nu_\mu$$

$$n \rightarrow p + e^- + \bar{\nu}_e$$

Resulting ν_μ flux (at the observer)



P. BAERWALD, S. HÜMMER, AND W. WINTER, *Phys. Rev. D* **83**, 067303 (2011)

NeuCosmA – how the neutrino spectrum changes – I

Corrections to the analytical model:

► **shape revised:**


- shift of first break (correction of photohadronic threshold)
- different cooling breaks for μ 's and π 's
- $(1+z)$ correction on the variability scale of the GRB

► **Correction cf_π to π prod. efficiency:**

- $f_{C\gamma}$: full spectral shape of photons
- $f_{\approx} = 0.69$: rounding error in analytical calculation
- $f_\sigma \simeq 2/3$: from neglecting the width of the Δ -resonance

► **Correction c_s :**


- energy losses of secondaries
- energy dependence of the mean free path of protons




S. HÜMMER, P. BAERWALD, W. WINTER,
Phys. Rev. Lett. 108, 231101 (2012)

NeuCosmA – neutrino spectra including flavor mixing

Electron neutrino spectrum



Muon neutrino spectrum



P. BAERWALD, S. HÜMMER, W. WINTER, *Phys. Rev. D* **83**, 067303 (2011)

Characteristic double peak structure from μ and π decay in both flavors,
additional peak from K^+ decay at 10^8 to 10^9 GeV

The need for km-scale neutrino telescopes

Expected ν flux from cosmological accelerators (Waxman & Bahcall 1997 & 1998):

$$E^2 \Phi_\nu \sim 10^{-8} \frac{f_\pi}{0.2} \left(\frac{\dot{\varepsilon}_{\text{CR}}^{[10^{10}, 10^{12}]} }{10^{44} \text{ erg Mpc}^{-3} \text{ yr}^{-1}} \right) \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

Integrated flux above 1 PeV:

$$\Phi_\nu (> 1 \text{ PeV}) \sim \int_{1 \text{ PeV}}^{\infty} \frac{10^{-8}}{E^2} dE \sim 10^{-20} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

Number of events from half of the sky (2π):

$$N_\nu \simeq 2\pi \cdot \Phi_\nu (> 1 \text{ PeV}) \cdot 1 \text{ yr} \cdot A_{\text{eff}} \approx (2.4 \times 10^{-10} \text{ cm}^{-2}) A_{\text{eff}} ,$$

where A_{eff} is the effective area of the detector

To detect $N_\nu > 1$ events per year, we need an area of

$$A_{\text{eff}} \gtrsim 0.4 \text{ km}^2$$

Therefore, we need km-scale detectors, like IceCube

Cosmogenic neutrinos

We have seen that protons interact with the cosmological photon fields (CMB, etc.), e.g.,


$$p + \gamma \rightarrow \Delta^+ \rightarrow \pi^+ + n ,$$

and neutrinos are created in the decays of the secondaries:

$$\begin{aligned}\pi^+ &\rightarrow \mu^+ + \nu_\mu \\ \mu^+ &\rightarrow \bar{\nu}_\mu + \nu_e + e^+ \\ n &\rightarrow p + e^- + \bar{\nu}_e\end{aligned}$$

These are called *cosmogenic neutrinos*

Cosmogenic neutrinos



P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Interaction with the photon backgrounds – I

- ▶ Energy loss rate (GeV s^{-1}):

$$b(E) \equiv \frac{dE}{dt}$$

- ▶ For pair production $p\gamma \rightarrow pe^+e^-$:

$$b_{e^+e^-}(E, z) = -\alpha r_0^2 (m_e c^2)^2 c \int_2^\infty d\xi n_\gamma \left(\frac{\xi m_e c^2}{2\gamma}, z \right) \frac{\phi(\xi)}{\xi^2}$$

- ▶ n_γ : isotropic photon background ($\text{GeV}^{-1} \text{ cm}^{-3}$)
- ▶ ξ : photon energy in units of $m_e c^2$
- ▶ proton energy: $E = \gamma m_p c^2$ ($\gamma \gg 1$)
- ▶ $\phi(\xi)$: (tabulated) integral in energy of outgoing e^-

G. BLUMENTHAL, *Phys. Rev. D* **1**, 1596 (1970)

H. BETHE, W. HEITLER, *Proc. Roy. Soc. A* **146**, 83 (1934)

Interaction with the photon backgrounds – II

Photohadronic interactions – $p\gamma$ interaction rate (s^{-1} per particle):

$$\Gamma_{p\gamma \rightarrow p'b}(E, z) = \frac{1}{2} \frac{m_p^2}{E^2} \int_{\frac{\epsilon_{\text{th}} m_p}{2E}}^{\infty} d\epsilon \frac{n_{\gamma}(\epsilon, z)}{\epsilon^2} \int_{\epsilon_{\text{th}}}^{2E\epsilon/m_p} d\epsilon_r \epsilon_r \sigma_{p\gamma \rightarrow p'b}^{\text{tot}}(\epsilon_r)$$

- 1 For given values of E and z , NeuCosmA calculates the cooling rate $t_{p\gamma}^{-1} \equiv -(1/E) b_{p\gamma}$ (s^{-1}) as


$$t_{p\gamma}^{-1}(E, z) = \sum_i^{\text{all channels}} \Gamma_{p \rightarrow p}^i(E, z) K^i,$$

with $K^i E$ the loss of energy per interaction

- 2 From this, we calculate back $b_{p\gamma}$ (GeV s^{-1}) ...
- 3 ... and the corresponding energy-loss term in the transport equation, $\partial_E(b_{p\gamma} Y_p)$.

Interaction lengths

Note that $L_{\text{CIB}} \gg L_{\text{CMB}}$:



Matches, e.g., [H. TAKAMI, K. MURASE, S. NAGATAKI, K. SATO, Astropart. Phys. 31, 201 \(2009\) \[0704.0979\]](#)

UHE ν 's in the GRB internal shock model

Secondary injection of neutrons, neutrinos ($\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}$)

$$Q'(E') = \int_{E'}^{\infty} \frac{dE'_p}{E'_p} N'_p(E'_p) \int_0^{\infty} c d\varepsilon' N'_\gamma(\varepsilon') R(E', E'_p, \varepsilon')$$

Normalisation to the observed GRB photon flux F_γ

$$\int \varepsilon' N'_\gamma(\varepsilon') d\varepsilon' = \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto F_\gamma, \quad \int E'_p N'_p(E'_p) dE'_p = \frac{1}{f_e} \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto \frac{F_\gamma}{f_e}$$

Fluence per shell, at Earth ($\text{GeV}^{-1} \text{ cm}^{-2}$)

$$\mathcal{F}^{\text{sh}} = t_\nu V'_{\text{iso}} \frac{(1+z)^2}{4\pi d_L^2} Q'$$

UHE ν 's in the GRB internal shock model

Secondary injection of neutrons, neutrinos ($\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}$)

$$Q' (E') = \int_{E'}^{\infty} \frac{dE'_p}{E'_p} \mathbf{N}'_p (E'_p) \int_0^{\infty} cd\varepsilon' \mathbf{N}'_{\gamma} (\varepsilon') R (E', E'_p, \varepsilon')$$

► Photon density, shock rest frame ($\text{GeV}^{-1} \text{ cm}^{-3}$):

$$\mathbf{N}'_{\gamma} (\varepsilon') \propto \begin{cases} (\varepsilon')^{-\alpha_{\gamma}}, & \varepsilon'_{\gamma, \text{min}} = 0.2 \text{ eV} \leq \varepsilon' \leq \varepsilon'_{\gamma, \text{break}} \\ (\varepsilon')^{-\beta_{\gamma}}, & \varepsilon'_{\gamma, \text{break}} \leq \varepsilon' \leq \varepsilon'_{\gamma, \text{max}} = 300 \times \varepsilon'_{\gamma, \text{min}} \end{cases}$$
$$\varepsilon'_{\gamma, \text{break}} = \mathcal{O}(\text{keV}), \alpha_{\gamma} \approx 1, \beta_{\gamma} \approx 2$$

► Proton density:

$$\mathbf{N}'_p (E'_p) \propto (E'_p)^{-\alpha_p} \times \exp \left[- \left(E'_p / E'_{p, \text{max}} \right)^2 \right] \quad (\alpha_p \approx 2)$$

Maximum proton energy limited by energy losses:

$$t'_{\text{acc}} (E'_{p, \text{max}}) = \min [t'_{\text{dyn}} (E'_{p, \text{max}}), t'_{\text{syn}} (E'_{p, \text{max}}), t'_{p\gamma} (E'_{p, \text{max}})]$$

UHE ν 's in the GRB internal shock model

Secondary injection of neutrons, neutrinos ($\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}$)

$$Q'(E') = \int_{E'}^{\infty} \frac{dE'_p}{E'_p} N'_p(E'_p) \int_0^{\infty} c d\varepsilon' N'_\gamma(\varepsilon') R(E', E'_p, \varepsilon')$$

Normalisation to the observed GRB photon flux F_γ

$$\int \varepsilon' N'_\gamma(\varepsilon') d\varepsilon' = \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto F_\gamma, \quad \int E'_p N'_p(E'_p) dE'_p = \frac{1}{f_e} \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto \frac{F_\gamma}{f_e}$$

UHE ν 's in the GRB internal shock model

Secondary injection of neutrons, neutrinos ($\text{GeV}^{-1} \text{ cm}^{-3} \text{ s}^{-1}$)

$$Q'(E') = \int_{E'}^{\infty} \frac{dE'_p}{E'_p} N'_p(E'_p) \int_0^{\infty} c d\varepsilon' N'_\gamma(\varepsilon') R(E', E'_p, \varepsilon')$$

Normalisation to the observed GRB photon flux F_γ



$$\int \varepsilon' N'_\gamma(\varepsilon') d\varepsilon' = \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto F_\gamma, \quad \int E'_p N'_p(E'_p) dE'_p = \frac{1}{f_e} \frac{E'_{\text{iso}}^{\text{sh}}}{V'_{\text{iso}}} \propto \frac{F_\gamma}{f_e}$$

Fluence per shell, at Earth ($\text{GeV}^{-1} \text{ cm}^{-2}$)

$$\mathcal{F}^{\text{sh}} = t_\nu V'_{\text{iso}} \frac{(1+z)^2}{4\pi d_L^2} Q'$$

Gamma-ray and neutrino pulses




A fast-rise-exponential-decay (FRED) gamma-ray pulse is emitted in every collision:






$$L_\gamma^{\text{peak}} \sim R_C^{-2}$$

The prediction *is* robust

Simulations show only weak dependence of the **flux** on the boost Γ ...




... and on the GRB engine variability time δt_{eng}



Accelerating iron

- ▶ Photodisintegration destroys nuclei close to the center ($\sim 10^8$ km)
e.g., [ANCHORDOQUI *et al.*, Astropart. Phys. 29, 1 \(2008\)](#)
- ▶ However, they can survive at large radii:



MB, BAERWALD, MURASE, WINTER
Nature Commun. 6, 6783 (2015)

Propagating the UHECRs to Earth

We use a **Boltzmann equation** to transport protons to Earth:

- ▶ Comoving number density of protons ($\text{GeV}^{-1} \text{ cm}^{-3}$):

$$Y_p(E, z) = n_p(E, z) / (1 + z)^3 ,$$

with n_p the real number density


- ▶ Transport equation (comoving source frame):

$$\dot{Y}_p = \underbrace{\partial_E (H E Y_p)}_{\text{adiabatic losses}} + \underbrace{\partial_E (b_{e^+ e^-} Y_p)}_{\text{pair production losses}} + \underbrace{\partial_E (b_{p\gamma} Y_p)}_{\text{photohadronic losses}} + \mathcal{L}_{\text{CR}}$$

$Q_{\text{CR}}(E) \propto E^{-\alpha_p} e^{-E/E_{p,\text{max}}}$

Contribution of GRBs to the diffuse ν flux

- ▶ **Three populations:** high-luminosity long GRBs (HL-GRB), low-luminosity long GRBs (LL-GRB), short GRBs (sGRB)
- ▶ **Sub-PeV:** GRBs contribute a few % to the IceCube diffuse flux
- ▶ **PeV:** contribution could be higher



I. TAMBORRA, S. ANDO, JCAP 1509, 036 (2015)


Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)

direct p escape, $\alpha_p = 2$, $\eta = 1.0$




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions
- 4 Find the baryonic loading (i.e., relative energy of p 's to e 's) at each point




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions
- 4 Find the baryonic loading (i.e., relative energy of p 's to e 's) at each point
- 5 Identify the region corresponding to pure n escape and to n escape + CR leakage




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every

direct p escape, $\alpha_p = 2$, $\eta = 1$




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:

- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions
- 4 Find the baryonic loading (i.e., relative energy of p 's to e 's) at each point
- 5 Identify the region corresponding to pure n escape and to n escape + CR leakage
- 6 Find the region where the number of prompt ν_μ 's is > 2.44 , i.e., the excluded region at 90% C.L.




P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Using UHECR + neutrino constraints

We can already limit the parameter space by using the UHECR observations and ν upper bounds:


- 1 Generate the UHECR spectrum at every point in parameter space (e.g., in Γ vs. L_{iso})
- 2 Fit each spectrum to UHECR data (TA, PAO, HiRes)
- 3 Find the best-fit point (diamond), and the 90% (red), 95% (yellow), and 99% (blue) C.L. regions
- 4 Find the baryonic loading (i.e., relative energy of p 's to e 's) at each point
- 5 Identify the region corresponding to pure n escape and to n escape + CR leakage
- 6 Find the region where the number of prompt ν_μ 's is > 2.44 , i.e., the excluded region at 90% C.L.
- 7 After 15 yr of exposure and no detection, cosmogenic neutrinos also exclude



P. BAERWALD, MB, W. WINTER, *Astropart. Phys.* **62**, 66 (2015)

Initialising the burst simulation


Initial number of plasma shells in the jet: $\gtrsim 1000$



Initial values of shell parameters:

- ▶ Width of shells and separation between them: $l = d = c \cdot \delta t_{\text{eng}}$
- ▶ Equal kinetic energy for all shells ($\sim 10^{52}$ erg)
- ▶ Shell speeds $\Gamma_{k,0}$ follow a distribution (log-normal or other)

Propagating and colliding the shells



During propagation:

- speeds, masses, widths **do not** change (only in collisions)
- the new, merged shells continue propagating and can collide again

Evolution stops when either:

- a single shell is left; or
- all remaining shells have reached the circumburst medium ($\gtrsim 6 \times 10^{11}$ km)

final number of collisions

\approx

number of initial shells ($\gtrsim 1000$)

S. KOBAYASHI, T. PIRAN, R. SARI, *ApJ* 490, 92 (1997)

F. DAIGNE, R. MOCHKOVITCH, *MNRAS* 296, 275 (1998)

How is the new prediction different?

- ▶ The top-contributing collisions are at the photosphere
- ▶ Pion production efficiency there is **independent of Γ** :

$$f_{p\gamma}^{\text{ph}} \sim 5 \cdot \frac{\varepsilon}{0.25} \cdot \frac{\epsilon_e}{0.1} \cdot \frac{1 \text{ keV}}{\epsilon'_{\gamma, \text{break}}}$$

ε : energy dissipation efficiency

ϵ_e : fraction of dissipated energy as e.m. output (photons)

- ▶ \Rightarrow Time-integrated neutrino fluence dominated is independent of Γ :

$$\mathcal{F}_\nu \propto \frac{N_{\text{coll}} (f_{p\gamma} \gtrsim 1)}{N_{\text{coll}}^{\text{tot}}} \times \min \left[1, f_{p\gamma}^{\text{ph}} \right] \times \frac{\epsilon_p}{\epsilon_e} \times E_{\gamma-\text{tot}}^{\text{iso}}$$

- ▶ Compare to standard predictions, which have a $\langle \Gamma \rangle^{-4}$ dependence
- ▶ Raising ϵ_p automatically decreases ϵ_e , so the photosphere grows, but still ~ 10 photospheric collisions dominate

How is the new prediction different?

- ▶ The top-contributing collisions are at the photosphere
- ▶ Pion production efficiency there is **independent of Γ** :

$$f_{p\gamma}^{\text{ph}} \sim 5 \cdot \frac{\varepsilon}{0.25} \cdot \frac{\epsilon_e}{0.1} \cdot \frac{1 \text{ keV}}{\epsilon'_{\gamma, \text{break}}}$$

ε : energy dissipation efficiency

ϵ_e : fraction of dissipated energy as e.m. output (photons)

- ▶ \Rightarrow Time-integrated neutrino fluence dominated is independent of Γ :

$$\mathcal{F}_\nu \propto \frac{\sim 10}{\frac{N_{\text{coll}}(f_{p\gamma} \gtrsim 1)}{N_{\text{coll}}^{\text{tot}}} \sim 1000} \times \min \left[1, f_{p\gamma}^{\text{ph}} \right] \times \frac{10}{\frac{\epsilon_p}{\epsilon_e}} \times 10^{53} \text{ erg} \times E_{\gamma-\text{tot}}^{\text{iso}}$$

- ▶ Compare to standard predictions, which have a $\langle \Gamma \rangle^{-4}$ dependence
- ▶ Raising ϵ_p automatically decreases ϵ_e , so the photosphere grows, but still ~ 10 photospheric collisions dominate