

High-energy neutrino physics from high-energy astrophysical neutrinos

Mauricio Bustamante

Niels Bohr Institute, University of Copenhagen


GRAPPA Colloquium
Amsterdam, December 09, 2024


UNIVERSITY OF
COPENHAGEN





VILLUM FONDEN

















They have the **highest energies**










Synergies with lower energies


Discovered in 2013
by IceCube



Synergies with lower energies







Today
TeV–PeV ν

Astro: Find & understand sources

Particle: Turn predictions into tests

Next decade
> 100-PeV ν

Make predictions for
a new energy regime

The story so far



Making high-energy astrophysical neutrinos: a toy model

(or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$


Making high-energy astrophysical neutrinos: a toy model

(or $p + p$)



Making high-energy astrophysical neutrinos: a toy model

(or $p + p$)



Proton density

Energy






$\sim E^{-2}$

Photon density

Energy

Making high-energy astrophysical neutrinos: a toy model

(or $p + p$)




Making high-energy astrophysical neutrinos: a toy model (or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \bar{\nu}_\mu + e^+ + \nu_e + \nu_\mu$$

$$n \text{ (escapes)} \rightarrow \textcolor{red}{p} + e^- + \bar{\nu}_e$$



Neutrino energy = Proton energy / 20


Gamma-ray energy = Proton energy / 10

Redshift



$z = 0$

Note: v sources can be steady-state or transient



Redshift

$z = 0$

Discovered

MeV γ

PeV p

TeV–PeV ν

“High-energy”


Photohadronic or pp interaction
inside the source

Note: ν sources can be steady-state or transient


ν propagation
inside the Earth

ν detection

HE ν




Today &
immediate
future




IceCube – What is it?

- ▶ Km^3 in-ice Cherenkov detector in Antarctica
- ▶ > 5000 PMTs at 1.5–2.5 km of depth
- ▶ Sensitive to neutrino energies > 10 GeV




Shower (mainly from ν_e and ν_τ)




Poor angular resolution: $\geq 5^\circ$

Track (mainly from ν_μ)



Angular resolution: $< 1^\circ$




Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)


Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ



Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)




Standard expectation:
 ν and γ from transients arrive simultaneously


Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Neutrino energy spectrum

With > 10 years of data, deviations from a power law start to be testable:



Different spectra might reflect different source populations



Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)




Standard expectation:
 ν and γ from transients arrive simultaneously


Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Distribution of arrival directions (7.5 yr)


No significant excess in the neutrino skymap:




High-energy neutrinos from the Galactic Plane




High-energy neutrinos from the Galactic Plane




High-energy neutrinos from the Galactic Plane



High-energy neutrinos from the Galactic Plane




High-energy neutrinos from the Galactic Plane



4.5 σ evidence (post-trial) of
diffuse flux of $> \text{TeV } \nu$ from the GP

Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)




Standard expectation:
 ν and γ from transients arrive simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S}) / N_{\text{tot}}$$


Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S}) / N_{\text{tot}}$$

Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$

Standard oscillations
or
new physics


Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




From sources to Earth: we learn what to expect when measuring $f_{\alpha,\oplus}$

Sources



E.g.,



$(f_{e,S}, f_{\mu,S}, f_{\tau,S})$

Oscillations

$(\theta_{12}, \theta_{23}, \theta_{13}, \delta_{CP})$

Earth



$(f_{e,\oplus}, f_{\mu,\oplus}, f_{\tau,\oplus})$

One likely TeV–PeV ν production scenario:


$$p + \gamma \rightarrow \pi^+ \rightarrow \mu^+ + \nu_{\mu} \text{ followed by } \mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_{\mu}$$

Full π decay chain

$$(1/3:2/3:0)_S$$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable
in neutrino telescopes

One likely TeV–PeV ν production scenario:




Full π decay chain

(1/3:2/3:0)_S

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

One likely TeV–PeV ν production scenario:



Full π decay chain

$(1/3:2/3:0)_S$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

$$p + \gamma \rightarrow \pi^+ \rightarrow \mu^+ + \textcolor{brown}{\nu}_\mu \quad \text{followed by} \quad \mu^+ \rightarrow e^+ + \textcolor{blue}{\nu}_e + \overline{\textcolor{brown}{\nu}_\mu}$$


$$(1/3:2/3:0)_S$$

- Muon damped

$$(0:1:0)_S$$

22

One likely TeV–PeV ν production scenario:



Full π decay chain

$(1/3:2/3:0)_S$

Muon damped

$(0:1:0)_S$


Neutron decay

$(1:0:0)_S$


Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

Measuring flavor composition: 2015–2020


Measuring flavor composition: 2015–2020



Measuring flavor composition: 2015–2020



Measuring flavor composition: 2015–2020




Measuring flavor composition: 2015–2020

Based on
real data


- 2015: IC, cont.+thr.
- 2018: IC, cont.
- 2020: IC, cont. (w/ ν_τ)

- π decay: $(1:2:0)_S$
- μ -damped: $(0:1:0)_S$
- n decay: $(1:0:0)_S$



Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)



Standard expectation:
 ν and γ from transients arrive simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ


Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays

$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right] \frac{f_p}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays



$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right] \frac{f_p}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays


$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right] \frac{f_p}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays


$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \underbrace{\left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle\right)^{\tau_{p\gamma}}\right]}_{\text{Fraction of total } p \text{ energy given to pions}} \frac{f_p}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays

Fraction of p energy given to π
in one interaction ($\sim 20\%$)

$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \underbrace{\left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right]}_{\text{Fraction of total } p \text{ energy given to pions}} \frac{f_p}{f_e} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays

Fraction of p energy given to π
in one interaction ($\sim 20\%$)

$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right] \left(\frac{f_p}{f_e} \right) \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Fraction of total p energy
given to pions

Baryonic loading

Bright in gamma rays, bright in high-energy neutrinos

Energy in neutrinos $\propto$ energy in gamma rays


Fraction of p energy given to π
in one interaction ($\sim 20\%$)

$$\int_0^\infty dE_\nu E_\nu F_\nu(E_\nu) = \frac{1}{8} \underbrace{\left[1 - \left(1 - \langle x_{p \rightarrow \pi} \rangle \right)^{\tau_{p\gamma}} \right]}_{\text{Fraction of total } p \text{ energy given to pions}} \underbrace{\left(\frac{f_p}{f_e} \right)}_{\text{Baryonic loading}} \int_{1 \text{ keV}}^{10 \text{ MeV}} dE_\gamma E_\gamma F_\gamma(E_\gamma)$$

Optical depth to $p\gamma$:
$$\tau_{p\gamma} = \left(\frac{L_\gamma^{\text{iso}}}{10^{52} \text{ ergs}^{-1}} \right) \left(\frac{0.01}{t_v} \right) \left(\frac{300}{\Gamma} \right)^4 \left(\frac{\text{MeV}}{\epsilon_{\gamma, \text{break}}} \right)$$

Gamma-ray bursts and blazars – *not* dominant

Gamma-ray bursts




Blazars




Gamma-ray bursts and blazars – *not* dominant

Gamma-ray bursts




1172 GRBs inspected, no correlation found
< 1% contribution to diffuse flux

Blazars



862 blazars inspected, no correlation found
< 27% contribution to diffuse flux


TXS 0506+056: The first *transient* source of high-energy ν



TXS 0506+056: The first *transient* source of high-energy ν

Blazar TXS 0506+056:

IceCube, *Science* 2018



After re-analysis (2101.09836),
significance dropped
from $p=7 \times 10^{-5}$ to $p=8 \times 10^{-3}$

2014–2015: 13 ± 5 ν flare, no X-ray flare
3.5 σ significance of correlation (post-trial)

2017: one 290-TeV ν + X-ray flare
1.4 σ significance of correlation

Combined (pre-trial): 4.1 σ

Today

TeV–PeV ν

Astro: Find & understand sources

Particle: Turn predictions into tests

Key developments:

Larger statistics

Better reconstruction

Smaller astrophysical uncertainties

High-energy neutrino physics *today*

Fundamental physics with high-energy cosmic neutrinos

- ▶ Numerous new ν physics effects grow as $\sim \kappa_n \cdot E^n \cdot L$
- ▶ So we can probe $\kappa_n \sim 4 \cdot 10^{-47} (E/\text{PeV})^{-n} (L/\text{Gpc})^{-1} \text{PeV}^{1-n}$
- ▶ Improvement over limits using atmospheric ν : $\kappa_0 < 10^{-29} \text{PeV}$, $\kappa_1 < 10^{-33}$

Fundamental physics with high-energy cosmic neutrinos


- ▶ Numerous new ν physics effects grow as $\sim \kappa_n \cdot E^n \cdot L$ $\left. \vphantom{\begin{matrix} E.g., \\ n = -1: \text{neutrino decay} \\ n = 0: \text{CPT-odd Lorentz violation} \\ n = +1: \text{CPT-even Lorentz violation} \end{matrix}} \right\} \begin{matrix} E.g., \\ n = -1: \text{neutrino decay} \\ n = 0: \text{CPT-odd Lorentz violation} \\ n = +1: \text{CPT-even Lorentz violation} \end{matrix}$
- ▶ So we can probe $\kappa_n \sim 4 \cdot 10^{-47} (E/\text{PeV})^{-n} (L/\text{Gpc})^{-1} \text{PeV}^{1-n}$
- ▶ Improvement over limits using atmospheric ν : $\kappa_0 < 10^{-29} \text{PeV}$, $\kappa_1 < 10^{-33}$




Note: Not an exhaustive list



Note: Not an exhaustive list



Note: Not an exhaustive list



Note: Not an exhaustive list


Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list




Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list




Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list




Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e, ν_μ, ν_τ

Note: Not an exhaustive list




Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list



More: *PoS ICRC2019* (1907.08690)

Argüelles, MB, Kheirandish, Palomares-Ruiz, Salvadó, Vincent


Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list



More: *PoS ICRC2019* (1907.08690)

Argüelles, MB, Kheirandish, Palomares-Ruiz, Salvadó, Vincent

Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Reviews:
Ahlers, Helbing, De los Heros, *EPJC* 2018
Argüelles, MB, Kheirandish, Palomares-Ruiz, Salvadó, Vincent, *ICRC* 2019 [1907.08690]
Ackermann, Ahlers, Anchordoqui, MB, et al., *Astro2020 Decadal Survey* [1903.04333]

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list

More: *PoS ICRC2019* (1907.08690)
Argüelles, MB, Kheirandish, Palomares-Ruiz, Salvadó, Vincent

A selection of neutrino physics

- 1 Discovering the Glashow resonance
 - 2 Neutrino-matter cross section
 - 3 Dark matter
 - 4 Flavor physics
 - 5 Secret neutrino interactions
 - 6 Neutrino decay
- } Find this in the backup slides


1. Glashow resonance:
Long-sought, finally seen

First observation of a Glashow resonance

Predicted in 1960:


First observation of a Glashow resonance

Predicted in 1960:





First observation of a Glashow resonance

Predicted in 1960:





First observation of a Glashow resonance

Predicted in 1960:




First observation of a Glashow resonance

Predicted in 1960:




First reported by IceCube in 2021:




First observation of a Glashow resonance

Predicted in 1960:




First reported by IceCube in 2021:




First observation of a Glashow resonance

Predicted in 1960:





First reported by IceCube in 2021:




First observation of a Glashow resonance

Predicted in 1960:





First reported by IceCube in 2021:




First observation of a Glashow resonance

Predicted in 1960:




First reported by IceCube in 2021:




2. Neutrino-matter cross section: *From TeV to PeV*

Center-of-mass energy $\sqrt{s}$ [GeV]




How does DIS probe nucleon structure?

What you see





Beneath the hood




(Plus the equivalent neutral-current process (Z-exchange))


Center-of-mass energy $\sqrt{s}$ [GeV]













Measuring the high-energy νN cross section

Below ~ 10 TeV: Earth is transparent




Above ~ 10 TeV: Earth is opaque




Measuring the high-energy νN cross section

Below ~ 10 TeV: Earth is transparent




Above ~ 10 TeV: Earth is opaque





Measuring the high-energy νN cross section

Below ~ 10 TeV: Earth is transparent




Above ~ 10 TeV: Earth is opaque







Center-of-mass energy $\sqrt{s}$ [GeV]




Center-of-mass energy $\sqrt{s}$ [GeV]



Center-of-mass energy $\sqrt{s}$ [GeV]



Center-of-mass energy $\sqrt{s}$ [GeV]



3. Dark matter:

Annihilation and decay into ν

High-energy neutrinos from dark matter

Dark matter co-annihilation:

$$\chi + \chi \rightarrow \nu + \bar{\nu}$$

$$\chi + \chi \rightarrow \dots \rightarrow \nu + \bar{\nu} + \dots$$

$$E_{\max} = m_{\chi}$$

Dark matter decay:


$$\chi \rightarrow \nu + \bar{\nu}$$

$$\chi \rightarrow \dots \rightarrow \nu + \bar{\nu} + \dots$$

$$E_{\max} = m_{\chi}/2$$

Electroweak corrections (off-shell W and Z emission) broaden the ν spectrum


$\nu + \bar{\nu}$ yield from DM (at source)



Approximate independence on m_{χ}
valid for $m_{\chi} \approx 100 \text{ TeV} - 10 \text{ PeV}$

Dark matter in the Milky Way



Fiorillo, Valera, MB, Winter, PRD 2023



DM annihilation
 $\Phi_\nu \propto \mathcal{I} \propto \rho_{\text{DM}}^2$


DM decay
 $\Phi_\nu \propto \mathcal{D} \propto \rho_{\text{DM}}$

IceCube, PRD 2023



Limits on dark matter annihilation


Per annihilation channel
(assuming 100% branching ratio)




Two DM contributions: Galactic (anisotropic) + extragalactic (isotropic)
Plus background of atmospheric neutrinos (anisotropic, but different)

Limits on dark matter annihilation

Per annihilation channel
(assuming 100% branching ratio)



Compared to other limits
(assuming annihilation to muons)



IceCube, JCAP 2023

Two DM contributions: Galactic (anisotropic) + extragalactic (isotropic)
Plus background of atmospheric neutrinos (anisotropic, but different)


4. New physics via flavor

Hard to do, but worth it

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S}) / N_{\text{tot}}$$


Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S}) / N_{\text{tot}}$$

Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$

Standard oscillations
or
new physics


Standard expectation:
Power-law energy spectrum

Standard expectation:
Isotropy (for diffuse flux)

Standard expectation:
 ν and γ from transients arrive
simultaneously

Standard expectation:
Equal number of ν_e , ν_μ , ν_τ

Note: Not an exhaustive list



More: *PoS ICRC2019* (1907.08690)

Argüelles, M.B., Kheirandish, Palomares-Ruiz, Salvadó, Vincent

New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

Reviews:


Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017

New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

► Neutrino decay

[Beacom *et al.*, *PRL* 2003; Baerwald, MB, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; MB, Beacom, Murase, *PRD* 2017]



Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017

New physics in flavor composition


Repurpose the flavor sensitivity to test new physics:

- Neutrino decay

[Beacom *et al.*, *PRL* 2003; Baerwald, **MB**, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; **MB**, Beacom, Murase, *PRD* 2017]

- Tests of unitarity at high energy

[Xu, He, Rodejohann, *JCAP* 2014; Ahlers, **MB**, Mu, *PRD* 2018;
Ahlers, **MB**, Nortvig, *JCAP* 2021]



Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017

New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

- Neutrino decay


[Beacom *et al.*, *PRL* 2003; Baerwald, **MB**, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; **MB**, Beacom, Murase, *PRD* 2017]

- Tests of unitarity at high energy

[Xu, He, Rodejohann, *JCAP* 2014; Ahlers, **MB**, Mu, *PRD* 2018;
Ahlers, **MB**, Nortvig, *JCAP* 2021]

- Lorentz- and CPT-invariance violation

[Barenboim & Quigg, *PRD* 2003; **MB**, Gago, Peña-Garay, *JHEP* 2010;
Kostelecky & Mewes 2004; Argüelles, Katori, Salvadó, *PRL* 2015]



Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017

New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

- ▶ Neutrino decay

[Beacom *et al.*, *PRL* 2003; Baerwald, **MB**, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; **MB**, Beacom, Murase, *PRD* 2017]

- ▶ Tests of unitarity at high energy

[Xu, He, Rodejohann, *JCAP* 2014; Ahlers, **MB**, Mu, *PRD* 2018;
Ahlers, **MB**, Nortvig, *JCAP* 2021]

- ▶ Lorentz- and CPT-invariance violation


[Barenboim & Quigg, *PRD* 2003; **MB**, Gago, Peña-Garay, *JHEP* 2010;
Kostelecky & Mewes 2004; Argüelles, Katori, Salvadó, *PRL* 2015]

- ▶ Non-standard interactions

[González-García *et al.*, *Astropart. Phys.* 2016;
Rasmussen *et al.*, *PRD* 2017]

Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017



New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

- ▶ Neutrino decay

[Beacom *et al.*, *PRL* 2003; Baerwald, **MB**, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; **MB**, Beacom, Murase, *PRD* 2017]

- ▶ Tests of unitarity at high energy

[Xu, He, Rodejohann, *JCAP* 2014; Ahlers, **MB**, Mu, *PRD* 2018;
Ahlers, **MB**, Nortvig, *JCAP* 2021]

- ▶ Lorentz- and CPT-invariance violation

[Barenboim & Quigg, *PRD* 2003; **MB**, Gago, Peña-Garay, *JHEP* 2010;
Kostelecky & Mewes 2004; Argüelles, Katori, Salvadó, *PRL* 2015]

- ▶ Non-standard interactions


[González-García *et al.*, *Astropart. Phys.* 2016;
Rasmussen *et al.*, *PRD* 2017]

- ▶ Active-sterile ν mixing

[Aeikens *et al.*, *JCAP* 2015; Brdar, Kopp, Wang, *JCAP* 2017;
Argüelles *et al.*, *JCAP* 2020; Ahlers, **MB**, *JCAP* 2021]

Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017



New physics in flavor composition

Repurpose the flavor sensitivity to test new physics:

- Neutrino decay

[Beacom *et al.*, *PRL* 2003; Baerwald, **MB**, Winter, *JCAP* 2010;
MB, Beacom, Winter, *PRL* 2015; **MB**, Beacom, Murase, *PRD* 2017]

- Tests of unitarity at high energy

[Xu, He, Rodejohann, *JCAP* 2014; Ahlers, **MB**, Mu, *PRD* 2018;
Ahlers, **MB**, Nortvig, *JCAP* 2021]

- Lorentz- and CPT-invariance violation

[Barenboim & Quigg, *PRD* 2003; **MB**, Gago, Peña-Garay, *JHEP* 2010;
Kostelecky & Mewes 2004; Argüelles, Katori, Salvadó, *PRL* 2015]

- Non-standard interactions

[González-García *et al.*, *Astropart. Phys.* 2016;
Rasmussen *et al.*, *PRD* 2017]

- Active-sterile ν mixing


[Aeikens *et al.*, *JCAP* 2015; Brdar, Kopp, Wang, *JCAP* 2017;
Argüelles *et al.*, *JCAP* 2020; Ahlers, **MB**, *JCAP* 2021]

- Long-range $e\nu$ interactions


[**MB** & Agarwalla, *PRL* 2019]

Reviews:

Mehta & Winter, *JCAP* 2011; Rasmussen *et al.*, *PRD* 2017




Lorentz-invariance violation can fill up the flavor triangle



$$H_{\text{tot}} = H_{\text{std}} + H_{\text{NP}}$$


$$H_{\text{std}} = \frac{1}{2E} U_{\text{PMNS}}^\dagger \text{diag} (0, \Delta m_{21}^2, \Delta m_{31}^2) U_{\text{PMNS}}$$

$$H_{\text{NP}} = \sum_n \left(\frac{E}{\Lambda_n} \right)^n U_n^\dagger \text{diag} (O_{n,1}, O_{n,2}, O_{n,3}) U_n$$




See also: Ahlers, **MB**, Mu, *PRD* 2018; Rasmusen *et al.*, *PRD* 2017; **MB**, Beacom, Winter *PRL* 2015; **MB**, Gago, Peña-Garay *JCAP* 2010; Bazo, **MB**, Gago, Miranda *IJMPA* 2009; + many others


Argüelles, Katori, Salvadó, *PRL* 2015



Dimension-5
CPT-odd
Isotropic
Lorentz-invariance
-violating
coefficient



High-energy neutrino physics *in the future*




The future

Build bigger

Build different

Work together




The future

Build bigger

Build different


Work together

Today &
immediate
future




Under construction


Early 2030s




Mid 2030s



Measuring the diffuse flux *precisely*




Assuming a power-law ν flux $\propto E^{-2.5}$




Assuming a power-law ν flux with 100-TeV cut-off + $p\gamma$ bump at tens of TeV


2040s




2040s




2040s



From today to
the 2040s







The future


Build bigger


Build different


Work together













Bump-hunting in the diffuse flux of high-energy neutrinos

Bump-like spectra can reveal the presence of ν production via $p\gamma$:




Bump-hunting in the diffuse flux of high-energy neutrinos

Bump-like spectra can reveal the presence of ν production via $p\gamma$:




Bump-hunting in the diffuse flux of high-energy neutrinos

Bump-like spectra can reveal the presence of ν production via $p\gamma$:




Bump-hunting in the diffuse flux of high-energy neutrinos

Bump-like spectra can reveal the presence of ν production via $p\gamma$:




Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021





Measuring flavor composition: 2015–2040

Song, Li, Argüelles, MB, Vincent, JCAP 2021



Measuring flavor composition: 2015–2040


Song, Li, Argüelles, MB, Vincent, JCAP 2021



How knowing the mixing parameters better helps


2020

NuFit 5.0




Measure θ_{23} better

+ Hyper-K




+ JUNO



Measure θ_{12} better

~2030

+ Hyper-K
+ JUNO




In our results:
JUNO + Hyper-K + DUNE


Marginal improvement til 2040


Theoretically palatable regions: 2020 $\rightarrow$ 2040

2020



2040







The future


Build bigger


Build different


Work together




















Redshift ←

| $z = 0$

MeV γ

Discovered

TeV–PeV ν

“High-energy”

PeV p


Photohadronic or pp interaction
inside the source

Note: ν sources can be steady-state or transient

HE ν

ν propagation
inside the Earth


ν detection



Redshift


$z = 0$

Note: ν sources can be steady-state or transient




Redshift ← $z = 0$


Note: ν sources can be steady-state or transient





Redshift ← $z = 0$


Note: ν sources can be steady-state or transient






















All-flavor neutrino flux, $E_\nu^2 \Phi_{\nu+\bar{\nu}}$ [$\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$]


- | | | |
|--|---|--|
| 1 IceCube HESE (7.5 yr) extrapolated | 4 Bergman & van Vliet, fit to TA UHECRs | 7 Rodrigues <i>et al.</i> , HL BL Lacs |
| 2 IceCube ν_μ (9.5 yr) extrapolated | 5 Rodrigues <i>et al.</i> , all AGN | 8 Fang & Murase, cosmic-ray reservoirs |
| 3 Heinze <i>et al.</i> , fit to Auger UHECRs | 6 Rodrigues <i>et al.</i> , all AGN | 9 Fang <i>et al.</i> , newborn pulsars |






All-flavor neutrino flux, $E_\nu^2 \Phi_{\nu+\bar{\nu}}$ [$\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$]

- | | | | |
|--|---|--|--|
| 1 IceCube HESE (7.5 yr) extrapolated | 4 Bergman & van Vliet, fit to TA UHECRs | 7 Rodrigues <i>et al.</i> , HL BL Lacs | 10 Padovani <i>et al.</i> , BL Lacs |
| 2 IceCube ν_μ (9.5 yr) extrapolated | 5 Rodrigues <i>et al.</i> , all AGN | 8 Fang & Murase, cosmic-ray reservoirs | 11 Muzio <i>et al.</i> , maximum extra p component |
| 3 Heinze <i>et al.</i> , fit to Auger UHECRs | 6 Rodrigues <i>et al.</i> , all AGN | 9 Fang <i>et al.</i> , newborn pulsars | |




All-flavor neutrino flux, $E_\nu^2 \Phi_{\nu+\bar{\nu}}$ [$\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$]

- | | | | |
|--|---|--|--|
| 1 IceCube HESE (7.5 yr) extrapolated | 4 Bergman & van Vliet, fit to TA UHECRs | 7 Rodrigues <i>et al.</i> , HL BL Lacs | 10 Padovani <i>et al.</i> , BL Lacs |
| 2 IceCube ν_μ (9.5 yr) extrapolated | 5 Rodrigues <i>et al.</i> , all AGN | 8 Fang & Murase, cosmic-ray reservoirs | 11 Muzio <i>et al.</i> , maximum extra p component |
| 3 Heinze <i>et al.</i> , fit to Auger UHECRs | 6 Rodrigues <i>et al.</i> , all AGN | 9 Fang <i>et al.</i> , newborn pulsars | 12 Muzio <i>et al.</i> , fit to Auger & IceCube |





All-flavor neutrino flux, $E_\nu^2 \Phi_{\nu+\bar{\nu}}$ [$\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$]




All-flavor neutrino flux, $E_\nu^2 \Phi_{\nu+\bar{\nu}}$ [$\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$]

- | | | | |
|--|---|--|--|
| 1 IceCube HESE (7.5 yr) extrapolated | 4 Bergman & van Vliet, fit to TA UHECRs | 7 Rodrigues <i>et al.</i> , HL BL Lacs | 10 Padovani <i>et al.</i> , BL Lacs |
| 2 IceCube ν_μ (9.5 yr) extrapolated | 5 Rodrigues <i>et al.</i> , all AGN | 8 Fang & Murase, cosmic-ray reservoirs | 11 Muzio <i>et al.</i> , maximum extra p component |
| 3 Heinze <i>et al.</i> , fit to Auger UHECRs | 6 Rodrigues <i>et al.</i> , all AGN | 9 Fang <i>et al.</i> , newborn pulsars | 12 Muzio <i>et al.</i> , fit to Auger & IceCube |







Center-of-mass energy $\sqrt{s}$ [GeV]



Center-of-mass energy $\sqrt{s}$ [GeV]



Center-of-mass energy $\sqrt{s}$ [GeV]



Today

TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Today
TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Next decade
 > 100 -PeV ν

Today
TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Next decade
 > 100 -PeV ν

Make predictions for
a new energy regime

Today
TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Next decade
 > 100 -PeV ν

Make predictions for
a new energy regime

Key developments:

Discovery

New detection techniques

Better UHE ν flux predictions

Today
TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Next decade
 > 100 -PeV ν

Make predictions for
a new energy regime

Key developments:

Discovery

New detection techniques

Better UHE ν flux predictions

Made robust and meaningful by accounting
for all relevant particle and astrophysics uncertainties

Today
TeV–PeV ν

Turn predictions
into data-driven tests

Key developments:

Bigger detectors $\rightarrow$ larger statistics

Better reconstruction

Smaller astrophysical uncertainties

Next decade
 $> 100\text{-PeV } \nu$

Make predictions for
a new energy regime

Key developments:

Discovery


New detection techniques

Better UHE ν flux predictions

Similar to the evolution of cosmology to a
high-precision field in the 1990s



Made robust and meaningful by accounting
for all relevant particle and astrophysics uncertainties



The future

Build bigger

Build different

Work together

How it
started

How it's
going

10–20 years
from now



How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν



How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered



How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

Hints of sources
First tests of ν physics

How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

Hints of sources
First tests of ν physics

EeV ν discovered
Precision tests with PeV ν
First tests with EeV ν

How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

Hints of sources
First tests of ν physics

How do we get there?

EeV ν discovered
Precision tests with PeV ν
First tests with EeV ν

Thanks!

IV PHD SUMMER SCHOOL ON NEUTRINOS

HERE, THERE & EVERYWHERE

NIELS BOHR INSTITUTE
COPENHAGEN • JULY 7–11, 2025

nbia.dk/neutrino2025

REGISTRATION DEADLINE: MARCH 31, 2025



For PhD and advanced MSc students • Organizers: Markus Ahlers & Mauricio Bustamante

UNIVERSITY OF
COPENHAGEN



The Niels Bohr
International Academy

VILLUM FONDEN




- ▶ One week
- ▶ Three tracks:
 - ▶ Neutrino **phenomenology**:
Mariam Tórtola (Valencia)
 - ▶ Neutrino **astrophysics**:
Maria Petropoulou (Athens)
 - ▶ Neutrino **cosmology**:
Vivian Poulin (Montepellier)
- ▶ Plus topical seminars & student talks
- ▶ Registration will open soon

nbia.dk/neutrino2025



Backup slides

Neutrino energy spectrum

7.5 yr: 100+ contained events > 60 TeV:




ν attenuated by Earth Atm. ν and μ vetoed




Data is fit well by a single power law:

$$\frac{d\Phi_{6\nu}}{dE_\nu} = \Phi_{\text{astro}} \left(\frac{E_\nu}{100 \text{ TeV}} \right)^{-\gamma_{\text{astro}}} \cdot 10^{-18} \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$




IceCube, ICRC 2023

High-energy neutrinos from the Galactic Plane



High-energy neutrinos from the Galactic Plane



Three models of Galactic diffuse ν :


π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV

Observed ($\times 5$ model)

Consistent with 100-TeV observations by Tibet Air Shower Array

Model

High-energy neutrinos from the Galactic Plane



Three models of Galactic diffuse ν :

π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV


KRA_γ^5 : Spectrum varies spatially, harder ν spectrum, cut-off at 5 PeV in CR energy

KRA_γ^{50} : Cut-off at 50 PeV in CR energy

Observed ($\times 0.5$ model)

Cut-off energy could be different from the 5 and 50 PeV tested

High-energy neutrinos from the Galactic Plane



Three models of Galactic diffuse ν :

π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV

KRA $_{\gamma}^5$: Spectrum varies spatially, harder ν spectrum, cut-off at 5 PeV in CR energy

KRA $_{\gamma}^{50}$: Cut-off at 50 PeV in CR energy

None of the models matched data

(caveat: there are relatively simple models)

No Galactic ν source identified

(likely diffuse + source: Fang & Murase, 2307.02905)

GP flux is 6–13% of all-sky at 30 TeV


NGC1068: The first *steady-state* source of high-energy ν


Active galactic nucleus

Brightest type-2 Seyfert


79^{+22}_{-20} ν of TeV energy

Significance: 4.2σ (global)






Neutrino-dark matter scattering



Signature:

Deficit of high-energy ν from the Galactic Center




21.3 $\log_{10}(\rho_{DM}/\text{GeV cm}^{-2})$ 23

7.5 years of IceCube HESE data

Upper limits (90% C.L.) on

$$g_{\text{max}} = \sqrt{g_\chi g_\nu}$$



IceCube, JCAP 2023
Argüelles, Kheirandish, Vincent, PRL 2017

Measuring the high-energy νN cross section

Number of detected neutrinos (simplified for presentation):

$$N \propto \underbrace{\Phi_\nu}_{\text{Neutrino flux}} \underbrace{\sigma_{\nu N}}_{\text{Cross section}} e^{-\tau_{\nu N}} = \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Measuring the high-energy νN cross section

Number of detected neutrinos (simplified for presentation):

$$N \propto \underbrace{\Phi_\nu}_{\text{Neutrino flux}} \underbrace{\sigma_{\nu N}}_{\text{Cross section}} e^{-\tau_{\nu N}} = \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Downgoing neutrinos
(L short $\rightarrow$ no matter)

$$N \propto \Phi_\nu \sigma_{\nu N}$$

Measuring the high-energy νN cross section

Number of detected neutrinos (simplified for presentation):

$$N \propto \underbrace{\Phi_\nu}_{\text{Neutrino flux}} \underbrace{\sigma_{\nu N}}_{\text{Cross section}} e^{-\tau_{\nu N}} = \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Downgoing neutrinos
(L short $\rightarrow$ no matter)

$$N \propto \underbrace{\Phi_\nu \sigma_{\nu N}}_{\text{Degeneracy}}$$

Measuring the high-energy νN cross section

Number of detected neutrinos (simplified for presentation):

$$N \propto \underbrace{\Phi_\nu}_{\text{Neutrino flux}} \underbrace{\sigma_{\nu N}}_{\text{Cross section}} e^{-\tau_{\nu N}} = \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Downgoing neutrinos
(L short $\rightarrow$ no matter)

$$N \propto \underbrace{\Phi_\nu \sigma_{\nu N}}_{\text{Degeneracy}}$$

Upgoing neutrinos
(L long $\rightarrow$ lots of matter)

$$N \propto \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Measuring the high-energy νN cross section

Number of detected neutrinos (simplified for presentation):

$$N \propto \underbrace{\Phi_\nu}_{\text{Neutrino flux}} \underbrace{\sigma_{\nu N}}_{\text{Cross section}} e^{-\tau_{\nu N}} = \Phi_\nu \sigma_{\nu N} e^{-L \sigma_{\nu N} n_N}$$

Downgoing neutrinos
(L short $\rightarrow$ no matter)

$$N \propto \underbrace{\Phi_\nu \sigma_{\nu N}}_{\text{Degeneracy}}$$


Upgoing neutrinos
(L long $\rightarrow$ lots of matter)

$$N \propto \Phi_\nu \sigma_{\nu N} \underbrace{e^{-L \sigma_{\nu N} n_N}}_{\text{Breaks the degeneracy}}$$

A feel for the in-Earth attenuation


Earth matter density

(Preliminary Reference Earth Model)





+


Neutrino-nucleon cross section




A feel for the in-Earth attenuation




No
attenuation




Full
attenuation






No
attenuation







Full
attenuation

















MB & Connolly, *PRL* 2019



MB & Connolly, *PRL* 2019




MB & Connolly, *PRL* 2019




MB & Connolly, *PRL* 2019

Peeking inside a proton




Limits on dark matter decay

Per annihilation channel
(assuming 100% branching ratio)




Compared to other limits
(assuming decay into muons)




Two DM contributions: Galactic (anisotropic) + extragalactic (isotropic)
Plus background of atmospheric neutrinos (anisotropic, but different)


No unitarity? No problem



vs.



How knowing the mixing parameters better helps



5. New neutrino interactions:
Are there secret $\nu\nu$ interactions?

Astrophysical neutrino sources

Earth




Galactic (kpc) or extragalactic (Mpc – Gpc) distance

Astrophysical neutrino sources

Earth

Galactic (kpc) or extragalactic (Mpc – Gpc) distance




Standard case: ν free-stream



(And oscillate)





Galactic (kpc) or extragalactic (Mpc – Gpc) distance





Galactic (kpc) or extragalactic (Mpc – Gpc) distance





Galactic (kpc) or extragalactic (Mpc – Gpc) distance





Galactic (kpc) or extragalactic (Mpc – Gpc) distance





Galactic (kpc) or extragalactic (Mpc – Gpc) distance



Galactic (kpc) or extragalactic (Mpc – Gpc) distance




Galactic (kpc) or extragalactic (Mpc – Gpc) distance



Secret interactions of high-energy astrophysical neutrinos

“Secret” neutrino interactions between astrophysical ν (PeV) and relic ν (0.1 meV):



Cross section:
$$\sigma = \frac{g^4}{4\pi} \frac{s}{(s - M^2)^2 + M^2\Gamma^2}$$

Resonance energy:
$$E_{\text{res}} = \frac{M^2}{2m_\nu}$$

MB, Rosenstroem, Shalgar, Tamborra, *PRD* 2020


See also: Esteban, Pandey, Brdar, Beacom, *PRD* 2021

Creque-Sarbinowski, Hyde, Kamionkowski, *PRD* 2021

Ng & Beacom, *PRD* 2014


Cherry, Friedland, Shoemaker, 1411.1071

Blum, Hook, Murase, 1408.3799



Secret interactions of high-energy astrophysical neutrinos

“Secret” neutrino interactions between astrophysical ν (PeV) and relic ν (0.1 meV):




Cross section:
$$\sigma = \frac{g^4 s}{4\pi (s - M^2)^2 + M^2\Gamma^2}$$

New coupling Mediator mass


Resonance energy:
$$E_{\text{res}} = \frac{M^2}{2m_\nu}$$

MB, Rosenstroem, Shalgar, Tamborra, *PRD* 2020
 See also: Esteban, Pandey, Brdar, Beacom, *PRD* 2021
 Creque-Sarbinowski, Hyde, Kamionkowski, *PRD* 2021
 Ng & Beacom, *PRD* 2014
 Cherry, Friedland, Shoemaker, 1411.1071
 Blum, Hook, Murase, 1408.3799



Secret interactions of high-energy astrophysical neutrinos

“Secret” neutrino interactions between astrophysical ν (PeV) and relic ν (0.1 meV):




Cross section:
$$\sigma = \frac{g^4 s}{4\pi (s - M^2)^2 + M^2\Gamma^2}$$

New coupling Mediator mass


Resonance energy:
$$E_{\text{res}} = \frac{M^2}{2m_\nu}$$

MB, Rosenstroem, Shalgar, Tamborra, *PRD* 2020
 See also: Esteban, Pandey, Brdar, Beacom, *PRD* 2021
 Creque-Sarbinowski, Hyde, Kamionkowski, *PRD* 2021
 Ng & Beacom, *PRD* 2014
 Cherry, Friedland, Shoemaker, 1411.1071
 Blum, Hook, Murase, 1408.3799



Secret interactions of high-energy astrophysical neutrinos

“Secret” neutrino interactions between astrophysical ν (PeV) and relic ν (0.1 meV):




Cross section:
$$\sigma = \frac{g^4}{4\pi} \frac{s}{(s - M^2)^2 + M^2\Gamma^2}$$

New coupling Mediator mass


Resonance energy:
$$E_{\text{res}} = \frac{M^2}{2m_\nu}$$

MB, Rosenstroem, Shalgar, Tamborra, *PRD* 2020
 See also: Esteban, Pandey, Brdar, Beacom, *PRD* 2021
 Creque-Sarbinowski, Hyde, Kamionkowski, *PRD* 2021
 Ng & Beacom, *PRD* 2014
 Cherry, Friedland, Shoemaker, 1411.1071
 Blum, Hook, Murase, 1408.3799



Secret interactions of high-energy astrophysical neutrinos

“Secret” neutrino interactions between astrophysical ν (PeV) and relic ν (0.1 meV):



Cross section:
$$\sigma = \frac{g^4 s}{4\pi (s - M^2)^2 + M^2 \Gamma^2}$$

New coupling g^4 (circled in red)

Mediator mass M^2 (circled in green)


Resonance energy:
$$E_{\text{res}} = \frac{M^2}{2m_\nu}$$

MB, Rosenstroem, Shalgar, Tamborra, *PRD* 2020
See also: Esteban, Pandey, Brdar, Beacom, *PRD* 2021
Creque-Sarbinowski, Hyde, Kamionkowski, *PRD* 2021
Ng & Beacom, *PRD* 2014
Cherry, Friedland, Shoemaker, 1411.1071
Blum, Hook, Murase, 1408.3799



Looking for evidence of ν SI

- ▶ Look for dips in 6 years of public IceCube data (HESE)
- ▶ 80 events, 18 TeV–2 PeV
- ▶ Assume flavor-diagonal and universal: $g_{\alpha\alpha} = g \delta_{\alpha\alpha}$
- ▶ Bayesian analysis varying M, g , shape of emitted flux (γ)
- ▶ Account for atmospheric ν , in-Earth propagation, detector uncertainties



No significant ($> 3\sigma$) evidence for a spectral dip ...



No significant ($> 3\sigma$) evidence for a spectral dip so we set upper limits on the coupling g



No significant ($> 3\sigma$) evidence for a spectral dip so we set upper limits on the coupling g



6. Unstable neutrinos: *Are neutrinos for ever?*

Are neutrinos forever?

- ▶ In the Standard Model (vSM), neutrinos are essentially stable ($\tau > 10^{36}$ yr):
 - ▶ One-photon decay ($\nu_i \rightarrow \nu_j + \gamma$): $\tau > 10^{36} (m_i/\text{eV})^{-5}$ yr
 - ▶ Two-photon decay ($\nu_i \rightarrow \nu_j + \gamma + \gamma$): $\tau > 10^{57} (m_i/\text{eV})^{-9}$ yr
 - ▶ Three-neutrino decay ($\nu_i \rightarrow \nu_j + \nu_k + \bar{\nu}_k$): $\tau > 10^{55} (m_i/\text{eV})^{-5}$ yr
- ▶ BSM decays may have significantly higher rates: $\nu_i \rightarrow \nu_j + \phi$
- ▶ We work in a model-independent way:
the nature of ϕ is unimportant if it is invisible to neutrino detectors

» Age of Universe
(~ 14.5 Gyr)

Are neutrinos forever?

- ▶ In the Standard Model (vSM), neutrinos are essentially stable ($\tau > 10^{36}$ yr):

- ▶ One-photon decay ($\nu_i \rightarrow \nu_j + \gamma$): $\tau > 10^{36} (m_i/\text{eV})^{-5}$ yr

- ▶ Two-photon decay ($\nu_i \rightarrow \nu_j + \gamma + \gamma$): $\tau > 10^{57} (m_i/\text{eV})^{-9}$ yr

- ▶ Three-neutrino decay ($\nu_i \rightarrow \nu_j + \nu_k + \bar{\nu}_k$): $\tau > 10^{55} (m_i/\text{eV})^{-5}$ yr

» Age of Universe
(~ 14.5 Gyr)

- ▶ BSM decays may have significantly higher rates: $\nu_i \rightarrow \nu_j + \phi$

Nambu-Goldstone
boson of a broken
symmetry


- ▶ We work in a model-independent way:

the nature of ϕ is unimportant if it is invisible to neutrino detectors

Astrophysical sources

Earth

$L \sim$ up to a few Gpc



The flux of ν_i is attenuated by $\exp[- (L/E) \cdot (\underbrace{m_i}_{\text{Mass of } \nu_i} / \underbrace{\tau_i}_{\text{Lifetime of } \nu_i})]$

Astrophysical sources

Earth

$L \sim$ up to a few Gpc



Decay changes the number
of each ν mass eigenstate, N_1, N_2, N_3

?



Only sensitive to their ratio

The flux of ν_i is attenuated by $\exp[- (L/E) \cdot \overbrace{(m_i/\tau_i)}^{\text{Mass of } \nu_i \text{ Lifetime of } \nu_i}]$

Astrophysical sources

Earth

$L \sim$ up to a few Gpc



Decay changes the number
of each ν mass eigenstate, N_1 , N_2 , N_3

?



Lower- E ν are longer-lived...


The flux of ν_i is attenuated by $\exp[- (L/E) \cdot (m_i/\tau_i)]$

... but ν that travel longer L are more attenuated!

Astrophysical sources

Earth

$L \sim$ up to a few Gpc



Astrophysical sources


Earth

$L \sim \text{up to a few Gpc}$

$$\nu_2, \nu_3 \rightarrow \nu_1$$

ν_1 lightest and stable
(normal mass ordering)

E.g.,




(If decay is complete)




Astrophysical sources


Earth

$L \sim \text{up to a few Gpc}$





$L \sim \text{up to a few Gpc}$



Fine print:


- Decay can be incomplete
- Final-state ν might be detectable or not
- Many more possible decay channels (see [Winter & Mehta, JCAP 2011](#))

What does neutrino decay change?


Flavor composition $\longleftrightarrow$ Spectrum shape $\longleftrightarrow$ Event rate

What does neutrino decay change?

Flavor composition



Spectrum shape



Event rate


Flavor content of mass eigenstates:

Known to within 2%

$$|U_{\alpha i}|^2 = |U_{\alpha i}(\theta_{12}, \theta_{23}, \theta_{13}, \delta_{\text{CP}})|^2$$

Known to within 8%

Known to within 20%
(or worse)



What does neutrino decay change?


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844


Flavor composition



Spectrum shape



Event rate



What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / MB, 2004.06844


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844



Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844



Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / **MB**, 2004.06844



Flavor composition



Spectrum shape



Event rate



What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / MB, 2004.06844 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape



Event rate




MB, Beacom, Murase, *PRD* 2017


What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape




Event rate




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape



Event rate



Glashow resonance (GR):

$\bar{\nu}_e + e \rightarrow W \rightarrow \text{hadrons} \rightarrow \text{shower}$

What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape




Event rate



Glashow resonance (GR):
 $\bar{\nu}_e + e \rightarrow W \rightarrow \text{hadrons} \rightarrow \text{shower}$

IceCube has seen one GR candidate in 4.6 years:




IceCube Collab., *Nature* 2021


What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape




Event rate



Glashow resonance (GR):

$$\bar{\nu}_e + e \rightarrow W \rightarrow \text{hadrons} \rightarrow \text{shower}$$


ν_1 is the mass eigenstate with the most e flavor




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape



Event rate



Glashow resonance (GR):


$\bar{\nu}_e + e \rightarrow W \rightarrow \text{hadrons} \rightarrow \text{shower}$

If $\bar{\nu}_1$ had decayed en route to Earth,
there would not have been $\bar{\nu}_e$ left to trigger a GR


What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, MB, Winter, *JCAP* 2012 / MB, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, MB, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape



Event rate



Glashow resonance (GR):

$\bar{\nu}_e + e \rightarrow W \rightarrow \text{hadrons} \rightarrow \text{shower}$


So by having observed 1 GR event we can place a *lower* limit on the lifetime of $\bar{\nu}_1$ ($= \nu_1$)

If $\bar{\nu}_1$ had decayed en route to Earth, there would not have been $\bar{\nu}_e$ left to trigger a GR


What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2020

Flavor composition




Spectrum shape



Event rate


MB, 2004.06844




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2020

Flavor composition




Spectrum shape



Event rate


MB, 2004.06844




What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape



Event rate




Free parameters:

ν_1, ν_2 lifetimes

Mix. par.

Flavor ratios

$\bar{\nu}/\nu$ ratio




MB, 2004.06844


What does neutrino decay change?

See also: Beacom *et al.*, *PRL* 2002 / Baerwald, **MB**, Winter, *JCAP* 2012 / **MB**, Beacom, Murase, *PRD* 2017 / Rasmussen *et al.*, *PRD* 2017 / Denton & Tamborra, *PRL* 2018 / Abdullahi & Denton, *PRD* 2020 / Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2020


Flavor composition



Spectrum shape





Event rate





Free parameters:


ν_1, ν_2 lifetimes
Mix. par.
Flavor ratios
 $\bar{\nu}/\nu$ ratio























① IceCube




IceCube + KM3NeT




PLE ν M-1 (① - ④)



PLE ν M-2 (② - ⑤)





PLE ν M-3 (② - ⑧)




↔ IceCube only

Rate of detected muon tracks relative to IceCube maximum







① IceCube




IceCube + KM3NeT




PLE ν M-1 (① - ④)



PLE ν M-2 (② - ⑤)




PLE ν M-3 (② - ⑧)




← IceCube only

Rate of detected muon tracks relative to IceCube maximum




Bump-hunting in the diffuse flux of high-energy neutrinos

Bump-like spectra can reveal the presence of ν production via $p\gamma$:




Radio emission: geomagnetic and Askaryan

Geomagnetic



- ▶ Time-varying transverse current
- ▶ Linearly polarized parallel to Lorentz force
- ▶ Dominant in air showers

Askaryan






- ▶ Time-varying negative-charge ~20% excess
- ▶ Linearly polarized towards axis
- ▶ Sub-dominant in air showers

Figures by H. Schoorlemmer and K. D. de Vries


Radio emission: geomagnetic and Askaryan

IceCube-Gen2 Radio






Giant Radio Array for Neutrino Detection



New physics in the UHE νN cross section

New physics in the UHE νN cross section


Heavy sterile neutrinos
via the dipole portal



Huang, Jana, Lindner, Rodejohann, 2204.10347


New physics in the UHE νN cross section

Heavy sterile neutrinos
via the dipole portal



Huang, Jana, Lindner, Rodejohann, 2204.10347


Multiple ν_τ -induced
bangs



Huang, EPJC 2022 [2207.02222]


New physics in the UHE νN cross section

Heavy sterile neutrinos
via the dipole portal




Huang, Jana, Lindner, Rodejohann, 2204.10347

Multiple ν_τ -induced
bangs



Huang, EPJC 2022 [2207.02222]

Leptoquarks,
charged Higgs, etc.




Huang, Jana, Lindner, Rodejohann, JCAP 2022 [2112.09476]

Flavor at ultra-high energies

Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:




Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:

indistinct detection of all flavors
by IceCube-Gen2 (radio)



Manufacturing UHE flavor sensitivity with two detectors


What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:

indistinct detection of all flavors
by IceCube-Gen2 (radio)

+

predominant detection of ν_τ
by GRAND



Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:


indistinct detection of all flavors
by IceCube-Gen2 (radio)

+

predominant detection of ν_τ
by GRAND

=

sensitivity to the fraction of UHE ν_τ



Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:


indistinct detection of all flavors
by IceCube-Gen2 (radio)

+

predominant detection of ν_τ
by GRAND

=

sensitivity to the fraction of UHE ν_τ



Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:


indistinct detection of all flavors
by IceCube-Gen2 (radio)

+

predominant detection of ν_τ
by GRAND

=

sensitivity to the fraction of UHE ν_τ



Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:


indistinct detection of all flavors
by IceCube-Gen2 (radio)

+

predominant detection of ν_τ
by GRAND

=

sensitivity to the fraction of UHE ν_τ



Manufacturing UHE flavor sensitivity with two detectors

What if future UHE radio-detection neutrino telescopes cannot see flavor?

Then we combine two of detectors:


indistinct detection of all flavors
by IceCube-Gen2 (radio)

+


predominant detection of ν_τ
by GRAND

=


sensitivity to the fraction of UHE ν_τ




Lorentz-invariance violation at ultra-high energies



Lorentz-invariance violation at ultra-high energies




Lorentz-invariance violation at ultra-high energies




Dimension of Lorentz invariance-violation (LIV) operator, d

Lorentz-invariance violation at ultra-high energies




Dimension of Lorentz invariance-violation (LIV) operator, d


Lorentz-invariance violation at ultra-high energies



IceCube-Gen2 (radio) alone might measure flavor




IceCube-Gen2 (radio) alone might measure flavor



Accessing the full UHE flavor information

IceCube-Gen2 (no flavor-id) + GRAND:

Access to ν_τ fraction



IceCube-Gen2 (with flavor-id):

Access to ν_e fraction and $\nu_\mu + \nu_\tau$ fraction

