

ICRC 2025

Neutrino Rapporteur

Mauricio Bustamante

Niels Bohr Institute, University of Copenhagen

39th International Cosmic Ray Conference
Geneva, July 24, 2025

UNIVERSITY OF
COPENHAGEN



VILLUM FONDEN



“Now this is not the end.
It is not even the beginning of the end.
But it is, perhaps,
the end of the beginning.”

W. Churchill, 1942

1 Increased ability to extract physical insight

Evolved from counting events to sophisticated analyses looking for subtle features

2 Intense theoretical interpretation work fueled by data

More data and higher precision help navigate the theory landscape

3 On our way to having multiple km-scale instruments

IceCube + KM3NeT + Baikal-GBD

4 We have broken into a new energy regime

KM3NeT detected the first ultra-high-energy neutrino

5 We have ambitious plans for the short- and mid-term

More, larger, different neutrino telescopes

A photograph of the IceCube observatory in Antarctica. The central structure is a blue, multi-level building with a complex network of metal stairs and walkways. It is flanked by two tall, cylindrical, silver-colored storage tanks. The entire facility is situated on a vast, flat, snow-covered landscape under a clear sky. The word "IceCube" is overlaid in a large, white, serif font across the center of the image.


IceCube

Near the South Pole

Volume $\sim 1 \text{ km}^3$, 1.5–2.5 km deep

Completed in 2011 (86 strings)

Discovered PeV ν in 2013



IceCube diffuse TeV–PeV ν flux

Using MESE (1 TeV–10 PeV) **[new flux points]**: V. Basu+ 985
Using contained + uncontained (DNNCascades, WIP): Z. Rechav+ 1154
Unfolding ν_μ spectrum **[new flux points, 11 yr]**: L. van Rootsellar 1199
PeV flux, combined data sets (WIP) : Y. Yildizci+ 1217
Using tau neutrino-induced cascades **[new flux fit, 11 yr]**: Z. Chen+ 1011

IceCube source searches

Transient sources, tracks + cascades (WIP): J. Carpio+ 1007
Flaring X-ray binaries, tracks +cascades 13 yr (WIP): A. Kochocki+ 1077
SN2023ixf **[new upper limits]**: A. Mand+ 1109
Real-time search for gravitational-wave coincidence: Z. Marka+ 1113
Infrared galaxies (WIP): T. Pernice+ 1145
Enhancements from machine learning (WIP): L. Seen+ 1169
GeV–PeV ν recurrent nova T Coronae Borealis (WIP): J. Thwaites+ 1200

IceCube Galactic ν flux

Segmented Galactic Plane (WIP): L. Neste+ 1130
Tracks (starting + thru) + cascades (WIP): J. Hellrung+ 1056 |
J. Osborn+ 1135 |
M. Thiesmeyer+ 1193
Improvement via summary statistics: O. Janik+ 1132

IceCube GeV ν counterparts to high-energy ν

First upper limits, short time-scales: C. Raab+ 1150
Prospects: P. A. S. Myhr+ 1174

IceCube flavor composition

Contained events > 1 TeV (MESE) **[new]**: A. Balagopal+ 983
High Energy Starting Events > 60 TeV (HESE) **[new 12 yr]**: N. Lad+ 1198

IceCube Extremely High Energy Events

Diffuse flux > 10 PeV **[new upper limit, 12.6 yr]**: M. Meier+ 1122
Improvements from machine learning: M. Nakos+ 1127

IceCube alerts

Improved real-time reconstruction (WIP): G. Sommani+ 1184
Real-time track alert catalog IceCat-2 (WIP): A. Zegarelli+ 1224

IceCube neutrino tomography of the Earth

Using ν from 500 GeV to 100 TeV, 11 yr (WIP): A. Y. Wen 1211

IceCube event selection

New DAQ to detect neutron echoes: K. Dutta+ 1030
Northern track selection using GCNN: P. Soldin+ 1183

IceCube detector studies and upgrades

Status of ice model: D. Chirkin+ 1013
Mapping ice stratigraphy: A. Eimer+ 1035


KM3NeT

A 3D rendering of the KM3NeT neutrino detector array. The image shows a vast grid of vertical strings of detector modules suspended in the deep sea. A prominent yellow cylindrical module is visible near the top center. On the right, a detailed view of a detector module is shown, featuring a spherical structure with multiple circular sensors. The background is a deep blue ocean with a bright light source on the horizon.

Astroparticle Research Mediterranean Sea

[3.5 km deep, Sicily, running since 2015]

Final configuration: $> 1 \text{ km}^3$, 230 strings, 4100+ optical modules • **Today:** 33 strings



KM3NeT highlight:
D. Dornic, July 21

Oscillation Research with Cosmics in the Abyss

2.5 km deep, Toulon, running since 2017

Final configuration: 1 km^3 , 115 strings, 2070 optical modules • **Today:** 28 lines

KM3NeT UHE neutrino

Observation: M. Lamoureux 1018

Source search: M. Marconi+ 1110

Cosmogenic origin: A. Condorelli+ 1111

Galactic origin: M. Breuhaus 997

Blazars: M. Bendahman+ 986 | M. Lincetto+ 1100

Global view: C. Argüelles+ 1117

GRB blastwave: P. A. Sevlé Myhr+ 1173

KM3NeT/ARCA: diffuse TeV–PeV ν flux

Today [new limit ARCA6-8-19-21]: V. Tsourapis+ 1196

Full detector: C. Karagiannis+ 1073

KM3NeT source searches

Point & extended [new limit ARCA6-8-19-21]: A. Hejboer+ 1124

With ANTARES [new limits ARCA6-8-19-21]: B. Caiiffi 1002

CMZ & Cygnus OB2: M. Benhassi+ 1039

Galactic gamma-ray sources: V. Parisi+ 1140

Blazar stacking [new limit ARCA6-8-19-21]: F. Carenini+ 1005 | M. R. Musone+ 1125

ULIRG stacking [new limit ARCA6-8-19-21]: A. De Benedittis+ 1023

Low energy, with ORCA: I. Del Rosso+ 1026

CCSNe: S. El Hedri 1037

Real-time alerts: F. Filippini+ 1038

Seyfert stacking [new limit ARCA6-8-19-21]: W. Idrissi Ibnsalih+ 1065

Microquasars: F. Magnani 1108

GRB stacking: J. Palacios González+ 1139

KM3NeT/ARCA: Galactic ν flux

Galactic Ridge [new limit ARCA6-8-19-21]: F. Filippini 1039

KM3NeT prompt atmospheric ν

Sensitivity: Kalaczyński+ 1039

KM3NeT/ORCA neutrino oscillations

First KM3NeT Bayesian: A. Lazo Pedrajas 1092

KM3NeT detector studies and upgrades

Control unit: C. Bozza 993

Junction box: E. Giorgio+ 1043

Online analysis system: M. Mastrodicasa+ 1115

ARCA string calibration: F. Badaracco+ 1116

ARCA time calibration: F. Benfenati Gualandi+ 1137

Acquisition systems: D. Real+ 1152 | D. Real+ 1153

ARCA acoustic positioning: S. Viola+ 1156

ORCA optical water properties: A. Romanov+ 1158



Baikal-GVD

In Lake Baikal, Russia

Long-running succession of
neutrino telescopes (since 1995)

Depths up to ~1.4 km


Array of clusters of 8 strings

Today:

0.7 km³, 14 clusters, 4212 OMs


Final configuration (in ~3 yr):

~1 km³, 90 strings, 6000+ OMs



In use


Single-cluster tracks



- ✓ Low energy threshold
- ✓ Optimal sensitivity to nearly vertical tracks
- ✓ 90% of recorded track events

In development


Multi-cluster tracks



- ✓ Moderately low energy threshold
- ✓ Optimal sensitivity to inclined tracks
- ✓ Best angular resolution

ν_μ CC


Single-cluster cascades



- ✓ High energy threshold
- ✓ Good energy resolution
- ✓ Relatively rare events

NC, ν_e ν_τ CC

Multi-cluster cascades



- ✓ Very high energy threshold
- ✓ Excellent energy resolution
- ✓ Very rare events

Baikal-GVD overview:
G. Safronov, July 19

Baikal-GVD: diffuse TeV–PeV ν flux

Using 6 yr cascades [**detection at 5.1σ**]: R. Dvornický+ 1031 [also 2507.01893]

Using tracks (work in progress): G. Safronov+ 1162

Baikal-GVD Galactic ν flux

Using 6 yr cascades, 2.5σ : B. B. Ulzutuev+ 1197

Baikal-GVD source searches

Point sources, using tracks [**new limits 5 yr tracks**]: G. Safronov+ 1162

Baikal-GVD UHE neutrinos

Diffuse flux [**new limit 6 yr cascades**]: M. N. Sorokovikov 1186

Baikal-GVD detector studies and upgrades

New positioning system: B. A. Shaybonov 1175

Optical water properties: B. A. Shaybonov 1176

ANTARES

A low-angle, upward-looking photograph of the ANTARES neutrino detector. The image shows a complex network of vertical stainless steel support poles. Several large, spherical photomultiplier tubes (PMTs) are visible, each encased in a black, dome-shaped protective housing. The PMTs are arranged in a vertical column, with the central one being the most prominent. The background is a dark, corrugated metal ceiling. The overall lighting is dim, with some reflections on the metallic surfaces.


40 km off of Toulon (ORCA site)

Volume $\sim 0.01 \text{ km}^3$, 2.5 km deep

Operated 2008–2022 (12 strings)

Provided invaluable experience
for later neutrino telescopes
(like AMANDA did in ice!)

Post-decommissioning, provided
PMTs for other experiments



ANTARES overview:
M. Spurio, July 17



ANTARES overview

20 years of ANTARES: M. Spurio+ 1187

ANTARES Galactic ν flux

Using tracks and cascades, full data set: T. Cartraud 1009

ANTARES source searches

Point-like and extended, full data set: F. Salesa Greus+ 1191

Neutrino flares, full data set: A. Sánchez Losa+ 1066

Gamma-ray bursts, full data set: M. Scarnera+ 1167

ANTARES BSM search

Magnetic monopoles, full data set: A. Eddymaoui+ 1033

One

The diffuse flux of TeV–PeV neutrinos

Two

Neutrinos from the Milky Way

Three

Neutrino flavor composition


Four

The first neutrino of ultra-high energy

Five

The shape of things to come

Probability to reach this section



One

The diffuse flux of
TeV–PeV neutrinos



The multi-messenger connection: a simple picture

(or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$



The multi-messenger connection: a simple picture

(or $p + p$)








The multi-messenger connection: a simple picture

(or $p + p$)



The multi-messenger connection: a simple picture

(or $p + p$)



The multi-messenger connection: a simple picture


(or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \bar{\nu}_\mu + e^+ + \nu_e + \nu_\mu$$

$$n \text{ (escapes)} \rightarrow \textcolor{red}{p} + e^- + \bar{\nu}_e$$



Neutrino energy = Proton energy / 20

Gamma-ray energy = Proton energy / 10

The multi-messenger connection: a simple picture


(or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \bar{\nu}_\mu + e^+ + \nu_e + \nu_\mu$$

$$n \text{ (escapes)} \rightarrow \textcolor{red}{p} + e^- + \bar{\nu}_e$$



1 PeV

20 PeV

Neutrino energy = Proton energy / 20

Gamma-ray energy = Proton energy / 10

The multi-messenger connection: a simple picture


(or $p + p$)

$$p + \gamma_{\text{target}} \rightarrow \Delta^+ \rightarrow \begin{cases} p + \pi^0, & \text{Br} = 2/3 \\ n + \pi^+, & \text{Br} = 1/3 \end{cases}$$

$$\pi^0 \rightarrow \gamma + \gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \bar{\nu}_\mu + e^+ + \nu_e + \nu_\mu$$

$$n \text{ (escapes)} \rightarrow \textcolor{red}{p} + e^- + \bar{\nu}_e$$



1 PeV

20 PeV

Neutrino energy = Proton energy / 20


Gamma-ray energy = Proton energy / 10

Redshift



$z = 0$

Note: ν sources can be steady-state or transient



Redshift

$z = 0$

Discovered

MeV γ

PeV p

TeV–PeV ν


“High-energy”

Photohadronic or pp interaction
inside the source

Note: ν sources can be steady-state or transient

ν propagation
inside the Earth


ν detection



Diffuse TeV–PeV ν flux: IceCube

V. Basu, A. Balagopal, A. Karle,
PoS(ICRC2025)985

1 – New all-flavor flux measurement at 1 TeV–10 PeV



11 yr of Medium Energy Starting Events (MESE)

Cascades (ν_e, ν_μ, ν_τ), tracks (ν_μ), double cascades (ν_τ)

Applied additional selection cuts to remove low-energy atmospheric muons

First time IceCube has resolved structure in the cosmic neutrino spectrum at $> 4\sigma$

Features of neutrino production?

Two source populations?



New physics? (E.g., dark-matter decay/annihilation)

SPL b.f.: $\Phi \propto E^{-\gamma}$ ($\gamma = 2.55$)


BPL b.f.: $\Phi \propto \begin{cases} E^{-\gamma_1}, & E < E_b \\ E^{-\gamma_2}, & E > E_b \end{cases}$ ($\gamma_1 = 1.72, \gamma_2 = 2.84, E_b = 33 \text{ TeV}$)

Diffuse TeV–PeV ν flux: IceCube


1 – New all-flavor flux measurement at 1 TeV–10 PeV



2 – New measurement using cascades at > 10 TeV



11 yr of cascade data



Cascades (ν_e, ν_μ, ν_τ) and double cascades (ν_τ)

Background to double-cascade search:


- ν_e charged-current cascades
- ν_e, ν_μ, ν_τ neutral-current cascades
- ν_μ starting tracks

Extra cuts to find double cascades (+ self-veto):

- total energy $> 10^{4.5}$ GeV
- inter-cascade length > 10 m
- energy asymmetry


Produce ν_τ -enriched sample with 90% ν_τ purity
(Great for flavor measurements, see later)

2 – New measurement using cascades at > 10 TeV



11 yr of cascade data

Cascades (ν_e, ν_μ, ν_τ) and double cascades (ν_τ)



Single-power-law (SPL) fit to data,

$$\Phi = \Phi_0 \times \left(\frac{E}{100 \text{ TeV}} \right)^{-\gamma},$$

agrees with previous results

Best-fit values:

$$\Phi_0 = 1.83 \pm 0.21$$

$$\gamma = 2.68 \pm 0.06$$

2 – New measurement using cascades at > 10 TeV

Tried also a broken-power-law fit:

$$\Phi = \Phi_0 \times \begin{cases} \left(\frac{E}{E_b}\right)^{-\gamma_1}, & E < E_b \\ \left(\frac{E}{E_b}\right)^{-\gamma_2}, & E \geq E_b \end{cases},$$


but the low-energy branch is only weakly constrained

Best-fit values:

$$\gamma_1 = 0.70_{-0.70}^{+1.05}$$

$$\gamma_2 = 2.83 \pm 0.12$$

$$E_b = 25.7_{-4.3}^{+9.0} \text{ TeV}$$



Diffuse TeV–PeV ν flux: Baikal-GVD

R. Dvornický (for Baikal-GVD)




PoS(ICRC2025)1031

See also [arXiv:2507.01893](https://arxiv.org/abs/2507.01893)

First independent observation of the diffuse ν flux

6 yr of cascade data (> 10 TeV): 2018–2024

Restricted to upgoing directions to mitigate atmospheric muon background



Diffuse TeV–PeV ν flux: Baikal-GVD

R. Dvornický (for Baikal-GVD)



PoS(ICRC2025)1031

See also [arXiv:2507.01893](https://arxiv.org/abs/2507.01893)

First independent observation of the diffuse ν flux

6 yr of cascade data (> 10 TeV): 2018–2024

Restricted to upgoing directions to mitigate atmospheric muon background



Diffuse TeV–PeV ν flux: Baikal-GVD

R. Dvornický (for Baikal-GVD)



PoS(ICRC2025)1031

See also [arXiv:2507.01893](https://arxiv.org/abs/2507.01893)

First independent observation of the diffuse ν flux

6 yr of cascade data (> 10 TeV): 2018–2024

Restricted to upgoing directions to mitigate atmospheric muon background




Diffuse TeV–PeV ν flux: Baikal-GVD

R. Dvornický (for Baikal-GVD)

PoS(ICRC2025)1031

See also [arXiv:2507.01893](https://arxiv.org/abs/2507.01893)

First independent observation of the diffuse ν flux



Single-power-law fit
to Baikal-GVD data

Baikal-GVD results compatible
(~90% C.L.) with IceCube HESE
(dominated by cascades > 60 TeV)

⚠ Tension with other IceCube results
(possible reasons:
systematics, ν sky anisotropies, etc.)

Fits to different
IceCube data sets



Diffuse TeV–PeV ν flux: Baikal-GVD

R. Dvornický (for Baikal-GVD)

PoS(ICRC2025)1031

See also [arXiv:2507.01893](https://arxiv.org/abs/2507.01893)


First independent observation of the diffuse ν flux



Diffuse TeV–PeV ν flux: KM3NeT

V. Tsourapis, E. Drakopoulou, L. Kalousis,
C. Markou, A. Sinopoulou, E. Tzamariudaki (for KM3NeT)
PoS(ICRC2025)1196

First upper limit from ARCA



640 days of ARCA6+8+19+21 (2021–2023)

Uses upgoing tracks to mitigate atmospheric muon background

Uses Bayesian approach (frequentist also available)

Best-fit values (single power law):

$$\Phi_0 = 3.03 \times 10^{-18} \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$$

$$\gamma = 3.0$$

Insufficient evidence for a measurement


Simulations – atm. μ : 45.8 • atm. ν : 1466.9 • cosmic ν : 140.7 vs. Data – 1780

Diffuse TeV–PeV ν flux: KM3NeT

V. Tsourapis, E. Drakopoulou, L. Kalousis,
C. Markou, A. Sinopoulou, E. Tzamariudaki (for KM3NeT)
PoS(ICRC2025)1196

First upper limit from ARCA

Place upper limit instead




Diffuse TeV–PeV ν flux: ANTARES


M. Spurio, S. Navas (for ANTARES)

PoS(ICRC2025)1187

See also *JCAP* 08, 038 (2024) [arXiv:2407.00328]


Upper limit using full legacy data from ANTARES





New in 2025


Thanks to
Aswathi Balagopal
for providing the
butterflies!



The three existing large neutrino telescopes are independently probing the diffuse flux of TeV–PeV cosmic ν !

Thanks to Aswathi Balagopal for providing the butterflies!

And IceCube is finding structure in the energy spectrum



New in 2025


The three existing large neutrino telescopes are independently probing the diffuse flux of TeV–PeV cosmic ν !

Thanks to Aswathi Balagopal for providing the butterflies!


Two

Neutrinos
from the Milky Way


Neutrinos from the Galaxy




Neutrinos from the Galaxy




Neutrinos from the Galaxy



Neutrinos from the Galaxy




Neutrinos from the Galaxy




4.5 σ evidence (post-trial) of
diffuse flux of $> \text{TeV } \nu$ from the GP

Neutrinos from the Galaxy: IceCube



Neutrinos from the Galaxy: IceCube



Three models of Galactic diffuse ν :


π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV

Observed ($\times 5$ model)

Consistent with 100-TeV observations by Tibet Air Shower Array

Model

Neutrinos from the Galaxy: IceCube



Three models of Galactic diffuse ν :

π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV


KRA $_\gamma^5$: Spectrum varies spatially, harder ν spectrum, cut-off at 5 PeV in CR energy

KRA $_\gamma^{50}$: Cut-off at 50 PeV in CR energy

Observed ($\times 0.5$ model)

Cut-off energy could be different from the 5 and 50 PeV tested

Neutrinos from the Galaxy: IceCube



Three models of Galactic diffuse ν :

π^0 : MeV–GeV π^0 template inferred from gamma rays extrapolated to TeV

KRA_{γ}^5 : Spectrum varies spatially, harder ν spectrum, cut-off at 5 PeV in CR energy

KRA_{γ}^{50} : Cut-off at 50 PeV in CR energy

None of the models matched data


(caveat: there are relatively simple models)

No Galactic ν source identified

(likely diffuse + source: Fang & Murase, 2307.02905)

GP flux is 6–13% of all-sky at 30 TeV

Improvements **without** template fitting



$\pm 8^\circ$ width in Galactic latitude
 $-40^\circ < \text{Galactic longitude} < 40^\circ$

Divide the Galactic Plane into 3 generic segments
[other segmentation schemes, too (e.g., 2, 6)]

Same cascade sample as 2023 discovery

Same unbinned maximum likelihood
... but now segmented


In each segment: single power law
Fit flux normalization and spectral index

Note: No systematics yet

Neutrinos from the Galaxy: IceCube

L. NESTE, M. Hünnefeld, C. Finley
PoS(ICRC2025)1130 • Talk July 21


Improvements **without** template fitting




Neutrinos from the Galaxy: IceCube

L. Nester, M. Hünnefeld, C. Finley
PoS(ICRC2025)1130 • Talk July 21

Improvements **without** template fitting




Improvements **without** template fitting



Neutrinos from the Galaxy: Baikal-GVD

Y. Y. Kovalev, A. V. Plavin, D. V. Semikoz,
V. Troitsky, B. B. Ulzutuev
PoS(ICRC2025)1197 • Talk July 19

Using 6 yr of cascades, 2.5σ



⊙ Detected cascade

Example:
20° band around the
Galactic Plane

Median Galactic latitude of events: $|b_{\text{med}}|$

If events are concentrated along the Plane,
 $|b_{\text{med}}|$ should be small


Method by [Y. Kovalev et al. ApJ 2022 \[2208.08423\]](#)
(formerly applied to IceCube tracks)

8 cascade events > 200 TeV, 2018–2024
~50% expected to be background

Neutrinos from the Galaxy: Baikal-GVD

Y. Y. Kovalev, A. V. Plavin, D. V. Semikoz,
V. Troitsky, B. B. Ulzutuev
PoS(ICRC2025)1197

Using 6 yr of cascades, 2.5σ



From data: $|b_{\text{med}}| = 10.4^\circ$

From simulations: $\langle |b_{\text{med}}| \rangle \approx 31^\circ$

p -value of observation: 0.014
(2.5σ significance, two-sided)

Rough flux estimate:

$$F_{\text{MW}} \propto \frac{n_{\text{MW}}}{n_{\text{all-sky}}} \Phi_{\text{all-sky}}$$

Num. events $|b_{\text{med}}| < 10^\circ (\approx 2.8)$ n_{MW} All-sky astro flux (uses Baikal flux)

Num. astro events all-sky (5.1) $n_{\text{all-sky}}$


$$\approx 20.1_{-0.9}^{+2.0} \times 10^{-18} \left(\frac{E}{100 \text{ TeV}} \right)^{-\gamma}$$

$(\gamma = 2.58_{-0.33}^{+0.27})$

Template-based search

Full ANTARES data set, 2007–2022, all-flavor: 3392 tracks + 406 showers (> 300 GeV)


No evidence found — instead 90% C.L. upper limits placed on multiple template models



Look for neutrinos from the Galactic Ridge

Galactic Ridge
 $|b| < 2^\circ$, $|l| < 30^\circ$


Hard cosmic-ray spectrum
High star-formation rate



Off region
Data-driven
background

See also:
ANTARES search, [PLB 2023 \[2212.11876\]](#)

Look for neutrinos from the Galactic Ridge



KM3NeT/ARCA6+8+19+21
Same data as for diffuse flux

Upgoing + horizontal tracks
Resolution $< 0.2^\circ$ at PeV

No evidence for Galactic
Ridge found

New 90% C.L. upper limit

See also:
ANTARES search, [PLB 2023 \[2212.11876\]](#)


Three

Neutrino
flavor composition

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S})/N_{\text{tot}}$$


Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$

Astrophysical sources

Earth

Up to a few Gpc



Different production mechanisms yield different flavor ratios:

$$(f_{e,S}, f_{\mu,S}, f_{\tau,S}) \equiv (N_{e,S}, N_{\mu,S}, N_{\tau,S})/N_{\text{tot}}$$

Flavor ratios at Earth ($\alpha = e, \mu, \tau$):

$$f_{\alpha,\oplus} = \sum_{\beta=e,\mu,\tau} P_{\nu_\beta \rightarrow \nu_\alpha} f_{\beta,S}$$


Standard oscillations
or
new physics

From sources to Earth: we learn what to expect when measuring $f_{\alpha,\oplus}$

Sources



E.g.,



$(f_{e,S}, f_{\mu,S}, f_{\tau,S})$

Oscillations

$(\theta_{12}, \theta_{23}, \theta_{13}, \delta_{CP})$

Earth



$(f_{e,\oplus}, f_{\mu,\oplus}, f_{\tau,\oplus})$

One likely TeV–PeV ν production scenario:


$$p + \gamma \rightarrow \pi^+ \rightarrow \mu^+ + \nu_\mu \text{ followed by } \mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$$

Full π decay chain

$$(1/3:2/3:0)_S$$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable
in neutrino telescopes

One likely TeV–PeV ν production scenario:




Full π decay chain

$(1/3:2/3:0)_S$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

One likely TeV–PeV ν production scenario:




Full π decay chain

$(1/3:2/3:0)_S$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

One likely TeV–PeV ν production scenario:



Full π decay chain


$(1/3:2/3:0)_S$

Muon damped

$(0:1:0)_S$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes

One likely TeV–PeV ν production scenario:



Full π decay chain

$(1/3:2/3:0)_S$

Muon damped


$(0:1:0)_S$

Neutron decay


$(1:0:0)_S$

Note: ν and $\bar{\nu}$ are (so far) indistinguishable in neutrino telescopes


Measuring flavor composition 2015–2025




Measuring flavor composition 2015–2025




Measuring flavor composition 2015–2025



Measuring flavor composition 2015–2025




Measuring flavor composition 2015–2025




Measuring flavor composition 2015–2025

Presented at
ICRC 2025



Measuring flavor composition 2015–2025



Four

The First Neutrino of
ultra-high energy

The international journal of science / 13 February 2025

nature

COSMIC CATCHER

Deep-sea telescope detects
neutrino with highest
energy ever recorded

Article

Observation of an ultra-high-energy cosmic neutrino with KM3NeT

KM3NeT Collab. *Nature* 638, 376 (2025)

One muon detected with 120^{+110}_{-60} PeV

The international journal of science / 13 February 2025

nature

COSMIC CATCHER

Deep-sea telescope detects
neutrino with highest
energy ever recorded

Article

Observation of an ultra-high-energy cosmic neutrino with KM3NeT

KM3NeT Collab. *Nature* 638, 376 (2025)

One muon detected with 120^{+110}_{-60} PeV

But is it due to a neutrino?

Yes! Direction points underground,
after traveling 150 km through Earth

Inferred neutrino energy: 220^{+570}_{-110} PeV

(Assuming E^{-2} spectrum)

nature

COSMIC CATCHER

Deep-sea telescope detects
neutrino with highest
energy ever recorded

Article

Observation of an ultra-high-energy cosmic neutrino with KM3NeT

KM3NeT Collab. *Nature* 638, 376 (2025)

One muon detected with 120^{+110}_{-60} PeV

But is it due to a neutrino?

Yes! Direction points underground,
after traveling 150 km through Earth

Inferred neutrino energy: 220^{+570}_{-110} PeV


(Assuming E^{-2} spectrum)

**RECORD
BREAKER**


Energy estimation

Overview:
M. Lamoureux, July 17

Muon energy is estimated from
the number of triggered PMTs



Stochastic losses consistent
with a single high-energy muon



EeV-scale atmospheric muon?

Muon bundle?

Atmospheric neutrino?

Unlikely: expectation is $< 10^{-4}$ per year


Unlikely: expectation is $< 10^{-3}$ per year

Unlikely: $(1-5) \times 10^{-5}$ per year

KM3NeT *vs.* IceCube & Auger

Joint fits:
J. Mauro, July 22

Diffuse flux of high-energy
astrophysical ν




KM3NeT vs. IceCube & Auger

Joint fits:
J. Mauro, July 22

UHE ν flux inferred from KM3NeT event
when considered by itself

Flux is above upper limits!




KM3NeT vs. IceCube & Auger

Joint fits:
J. Mauro, July 22

UHE ν flux inferred from KM3NeT event
when considered by itself


Flux is above upper limits!





KM3NeT vs. IceCube & Auger

Joint fits:
J. Mauro, July 22


UHE ν flux inferred when considering
non-observation by IceCube & Auger






Redshift ← $z = 0$



Redshift ← $z = 0$



Redshift ← $z = 0$




Was it a cosmogenic neutrino?

Cosmogenic origin:
A. Condorelli, July 22

Assume population of nondescript, identical UHECR sources

UHECR flux fit to Auger spectrum + mass composition, **source abundance** $(1+z)^m$





Joint neutrino + cosmic-ray interpretation


Muzio, Yuan, Lu, arXiv:2502.06944


Joint fits ν + UHECRs:
T. Yuan, July 22


Joint fit to IceCube and KM3NeT ν data + Auger UHECR data
(Assuming 100 PeV for KM3-230213A)

















Where did it come from?

From the Southern Hemisphere
(RA = 94.3°, dec = -7.8°)


Not far from Milky Way plane
But likely **not** of Milky-Way origin

KM3NeT Collab. [arXiv:2502.08387](#)

Likely extragalactic origin

Few extragalactic sources
(blazars) near event position,
but no strong association

KM3NeT Collab. [arXiv:2502.08484](#)



Where did it come from?

From the Southern Hemisphere
(RA = 94.3°, dec = -7.8°)


Not far from Milky Way plane
But likely **not** of Milky-Way origin

KM3NeT Collab. [arXiv:2502.08387](#)

Likely extragalactic origin

Few extragalactic sources
(blazars) near event position,
but no strong association

KM3NeT Collab. [arXiv:2502.08484](#)




Where did it come from?

Compatible w/ diffuse blazar population
M. Bendahman, July 22

Used to constrain ν flux from GRBs
P. A. S. Myhr, July 21

Could be a year-long transient source

Li, Machado, Naredo-Tuero, Schwemberger, arXiv:2502.04508
Neronov, Oikonomou, Semikoz, arXiv:2502.12986



Beyond the Standard Model

New energies represent new opportunities to look for BSM physics, *e.g.*,

- UHE ν from decay of super-heavy dark matter

- UHE ν from primordial black holes

- Sterile-active ν transitions

- Lorentz-invariance violation

Caveat emptor!


Being able to explain KM3-230213A with BSM physics *does not* mean that a BSM explanation is preferred (always compute your Bayes factors!)

See [backup slides](#) for BSM proposals inspired by KM3-230213A

New upper limits on UHE neutrinos: IceCube

Search for UHE ν updated from 9 to 12.6 yr

New limit:
M. Meier, July 17



Strongest UHE limit today

Improvement due to 40% higher $\nu_{\mu} A_{\text{eff}}$:
Improved angular resolution
Looser muon bundle cuts

Repeating the joint fit of the UHE
KM3-230213A with IceCube and Auger
increases the tension from 2.5σ to 2.9σ


Also: new limits on UHECR proton fraction

New upper limits on UHE neutrinos: Auger and TA

Auger surface detectors 7 yr

Search for downgoing low (DGL) showers: $60^\circ < \theta < 75^\circ$

Improvement in triggers




Auger limit:
S. Sehgal, July 17

TA surface detectors 16 yr

Search for showers with $65^\circ < \theta < 85^\circ$

Improvement in reconstruction method



TA limit:
K. Takahashi July 21


New upper limits on UHE neutrinos: Baikal-GVD

First UHE ν flux limit placed by Baikal-GVD

New limit:


M. Sorokovikov, July 17

Matches IceCube and Auger limits at low energies



Five

The shape of
things to come



Astrophysical neutrino flux

How much do we know about them?

GeV

TeV


PeV


EeV

ZeV

Neutrino energy ($\propto$ probability of neutrino interaction)








The future

Build bigger

Build different

Work together



IceCube Upgrade

Seven new strings


Enhance sensitivity to $O(10 \text{ GeV}) \nu$

Deployment in austral summer 2025/26

Major goal: improve calibration in the whole detector (ice properties)

Multiple calibration devices to be installed

But also science!




New, multi-PMT digital optical modules




IceCube Collaboration

S. Partenheimer+ 78 | M. Vereecken+ 108 | J. Braun+ 448 | K. Dutta+ 1029 | A. Eimer+ 1034 | N. Chau+ 1037 | S. Fukami 1040 | S. Griffin+ 1044 | C. Rott+ 1069 | Y. Kobayashi 1076

Sensitivity to GeV ν transients




WIMP DM annihilation in the Sun




IceCube Collaboration

S. Partenheimer+ 78 | M. Vereecken+ 108 | J. Braun+ 448 | K. Dutta+ 1029 | A. Eimer+ 1034 |
N. Chau+ 1037 | S. Fukami 1040 | S. Griffin+ 1044 | C. Rott+ 1069 | Y. Kobayashi 1076



P-ONE: Pacific Ocean Neutrino Explorer



P-ONE-1 talk:
C. Lagunas, July 16

Next phase: P-ONE-1

First string of P-ONE

2025: Construction and testing

2026: Deployment

P-ONE Collaboration

V. Gousy-Leblanc 1047 | D. Ghuman+ 1042 |

C. Lagunas Gualda 1087, 1088 | V. A Parrish 1141 | B. Veenstra 1202

P-ONE: Pacific Ocean Neutrino Explorer

P-ONE-1 talk:
C. Lagunas, July 16

Cascadia Basin, Canada

Two pathfinder arrays

STRAW (2018) + **STRAW-b** (2020)

Measure water properties

Next phase: P-ONE-1

First string of P-ONE

2025: Construction and testing


2026: Deployment

P-ONE Collaboration

V. Gousy-Leblanc 1047 | D. Ghuman+ 1042 |

C. Lagunas Gualda 1087, 1088 | V. A Parrish 1141 | B. Veenstra 1202

Effective volume: $> 1 \text{ km}^3$
70 strings, 1400 optical modules



P-ONE: Pacific Ocean Neutrino Explorer

P-ONE-1 talk:
C. Lagunas, July 16

Cascadia Basin, Canada

Two pathfinder arrays

STRAW (2018) + **STRAW-b** (2020)

Measure water properties

Next phase: P-ONE-1

First string of P-ONE

2025: Construction and testing

2026: Deployment



P-ONE Collaboration


V. Gousy-Leblanc 1047 | D. Ghuman+ 1042 |

C. Lagunas Gualda 1087, 1088 | V. A Parrish 1141 | B. Veenstra 1202



DETECTORS

SURFACE • RADIO • OPTICAL



Cosmic Ray Surface Array


An air shower array that sits on top of the optical array
One surface station installed above each optical string



IceCube-Gen2 Optical Module

4x the sensitivity of IceCube's modules
9,600 new optical modules in total to be deployed in the ice

80 modules on each string, spaced 17 meters apart



High-energy
IceCube successor

Optical array:
TeV–PeV ν
 $\sim 8 \text{ km}^3$ ($8 \times$ IceCube)
206 strings
9.6k new DOMs

$\sim 10 \times$ rate of PeV ν
vs. IceCube


IceCube-Gen2 overview:
M. Kowalski, July 21

IceCube-Gen2 Collaboration


List of contributions: 2507.08667

IceCube-Gen2

IceCube




IceCube - 1 km³
Energy > 10¹¹ eV




Optical - 8 km³
Energy > 10¹² eV


Precise measurement of the TeV–PeV diffuse ν flux



Discovery of TeV–PeV ν transients (*à la* TXS 0506+056)



Source and particle-physics tests via neutrino flavor



TRIDENT: The tRopIcal Deep sea Neutrino Telescope

South China Sea • Effective volume: 8 km^3 • 1000 strings, 20,000 optical modules

Sensitivity to TeV–EeV ν

Uneven Penrose tiling array
Larger array with broader energy range

TRIDENT overview + flavor id.:
W. Tian, July 19



NEON: Neutrino Observatory in the Nanhai

South China Sea


Effective volume: 10 km^3

1200 strings, 21,600 optical modules



$\sim 100\text{m}$

40m





High-Energy Underwater Neutrino Telescope

South China Sea

Effective volume: 30 km^3



2304 strings, 55,296 optical modules

Sensitive to ν of 100 TeV–100 PeV

HUNT overview:
T. Q. Huang, July 21

HUNT Collaboration

T. Q. Huang 1061 | Z. K Zeng+ 1126 | C. Liu



9 clusters of 128 strings each

Muon track angular resolution as good as 0.05°
(for tracks of 6 km in length)



High-Energy Underwater Neutrino Telescope

HUNT Collaboration

T. Q. Huang 1061 | Z. K Zeng+ 1126 | C. Liu

South China Sea

Effective volume: 30 km^3

2304 strings, 55,296 optical modules

Sensitive to ν of 100 TeV–100 PeV


March 2024: Test string in Lake Baikal

January 2025: Test string in South China Sea

2025–2026: 7 strings, South China Sea (*Seastar*)


HUNT overview:


T. Q. Huang, July 21



9 clusters of 128 strings each

Muon track angular resolution as good as 0.05°
(for tracks of 6 km in length)








TAMBO: the Tau Air-Shower Mountain-Based Observatory

Today

TAMBO-4



Small test array
Electronics, detector
prototypes

2025-2028


TAMBO-100
(aka TAMBITO)



Pilot array
Deployment techniques,
background, and social science

2028 - ...

TAMBO-5k




Array capable of detecting O(1)
neutrinos per year in the PeV to
100 PeV energy range

TAMBO Collaboration

W. Thompson, 1194 | P. Zhelnin+ 981 | C. Argüelles+ 1091

TAMBO overview:
W. Thompson, July 16

TAMBO: the Tau Air-Shower Mountain-Based Observatory



TAMBO Collaboration

W. Thompson, 1194 | P. Zhelnin+ 981 | C. Argüelles+ 1091


TAMBO overview:
W. Thompson, July 16

Trinity

Atmospheric Cherenkov imaging applied to PeV neutrinos

Pioneered by MAGIC (pointing at Atlantic), ASHRA, and NTA (Mauna Kea)

Trinity: 3 arrays each of 6 mirrors of 10 m²




Trinity Collaboration

S. Stepanoff 1188 | D. A. Raudales+ 1136



Trinity Demonstrator: 1.36 m^2
Frisko Peak, Utah, deployed 2023

Trinity One: first telescope of the array
Sensitivity to neutrino fluxes






Demonstrator Funding

2021


2023

Demonstrator First Light



Trinity One Funding

2026





2031

Trinity Observatory Funding


Trinity Collaboration

S. Stepanoff 1188 | D. A. Raudales+ 1136






TeV–PeV:




Earth is *almost fully* opaque,
some upgoing ν still make it through

TeV–PeV:



Earth is *almost fully* opaque,
some upgoing ν still make it through

> 100 PeV:




Earth is *completely* opaque,
but horizontal ν still make it through


ARA: Askaryan Radio Array

Running since 2011

Proof of concept for larger arrays
(RNO-G, IceCube-Gen2)



Five radio stations at > 200 m underground




ARA Collaboration

S. Ali+ 977 | A. Bishop+ 991 | P. Dasgupta+ 1022 | P. Giri+ | M. Muzio+ 1126 | M. F. H. Seikh+ 1171 | P. Windischhofer+ 1213 | A. Machtay | I. Kravchenko+


ARA: Askaryan Radio Array

First station-wide search for UHE ν



M. Muzio, July 17, PoS(ICRC2025)1126

First UHE ν search phased + traditional stations




P. Dasgupta, July 15, PoS(ICRC2025)1022

RNO-G: Radio Neutrino Observatory - Greenland


In construction

35 in-ice radio stations planned

ARA + ARIANNA experience






RNO-G overview:
A. Nelles, July 16



- 8 stations operational (Apr - Oct), Wind-power tested
- 4 additional stations drilled (as of today)

RNO-G: Radio Neutrino Observatory - Greenland

Flights over Antarctica
[[2506.17522](#)] and solar
flares [[2404.14995](#)] detected
(good for calibration)



Cosmic-ray searches
will be proof-of-concept
for neutrino searches

[A. Vijai, PoS\(ICRC2025\)1204 July 23](#)

[B. Hendricks, PoS\(ICRC2025\)1057](#)



RNO-G Collaboration

A. Nelles+ | F. Schüller+ 1168 | J. Henrichs+ 288 | B. Clark+ 1204 | B. Hendricks+ 1057 | M. Liu+ 319 | M. Punsuebsay 366 |
C. McLennan+ 288 | R. Camphyn 288 | J. Stoffels 1189 | M. Hossain Seikh 1356 | S. Bouma 135

RNO-G: Radio Neutrino Observatory - Greenland

Flight
[2500
flares [240
(good

Not drawn to scale




by searches
of-of-concept
o searches

RC2025)1204 July 23

oS(ICRC2025)1057

RNO-G Collaboration

A. Nelles+ | F. Schüller+ 116
C. McLennan+ 288 | R. Camp



Cosmic Ray Surface Array


An air shower array that sits on top of the optical array
One surface station installed above each optical string



IceCube-Gen2 Optical Module

4x the sensitivity of IceCube's modules
9,600 new optical modules in total to be deployed in the ice

80 modules on each string, spaced 17 meters apart



High-energy
IceCube successor

Radio array:
 $> 100 \text{ PeV } \nu$
 Askaryan radiation
 ~ 310 stations
 $\sim 500 \text{ km}^2$


$\sim 100\times$ rate of $\text{EeV } \nu$
vs. IceCube

IceCube-Gen2 radio:
 C. Glaser, July 21


IceCube-Gen2 Collaboration
 List of contributions: 2507.08667

IceCube-Gen2


IceCube



IceCube - 1 km³
Energy > 10¹¹ eV




Optical - 8 km³
Energy > 10¹² eV




Radio - 500 km³
Energy > 10¹⁶ eV

IceCube-Gen2 in-ice radio array


Detect the diffuse UHE ν flux even if it is small




Center-of-mass energy $\sqrt{s}$ [GeV]




Center-of-mass energy $\sqrt{s}$ [GeV]



IceCube-Gen2 in-ice radio array




RET: Radar Echo Telescope



Detect the ionization trail made by in-ice high-energy particle showers via active radar sounding

Today: RET-CR to detect cosmic rays, proof of principle


Future: RET-N to detect neutrinos



RET Collaboration

J. Loonen | D. Frikken | I. Loudon | K. N. Gopinath | A. Kyriacou

GRAND: Giant Radio Array for Neutrino Detection



Large array of antennas (50–200 MHz)

Detects geomagnetic radio emission from UHE particles

Three prototype arrays running since 2023:

GRANDProto300
65 antennas, China

• GRAND@Auger
10 antennas, Argentina

• GRAND@Nançay
5 antennas, France


GRAND overview:
O. Martineau, July 16

First cosmic-ray candidates shown at ICRC 2025

GRAND Collaboration

J. R. T. de Mello Neto 1024 | O. Martineau 1114 | P. Ma 453 | L. Piotrowski

GRAND: Giant Radio Array for Neutrino Detection



UHE ν sensitivity
reachable with full
detector (~200k
antennas, distributed)


Sufficient to test
UHE diffuse flux models
and transients (not shown)

Next phase: GRAND10k

HERON: Hybrid Elevated Radio Observatory for Neutrinos

BEACON phased-array stations (low-SNR detection) + GRAND antennas (self-triggering)

HERON overview:
K. Kotera, July 16



GRAND + BEACON Collaborations

K. Kotera+1078 | A. Zeolla 1227

GRAND-type:
standalone antennas
15 around each station
autonomous & external triggering
reconstruction



BEACON-type:
phased stations
24 antennas per station
low energy threshold for triggering



HERON: Hybrid Elevated Radio Observatory for Neutrinos

BEACON phased-array stations (low-SNR detection) + GRAND antennas (self-triggering)


Narrow field of view, high sensitivity:

Phase 1: 24 BEACON stations + 360 GRAND antennas [$\approx$ GRAND10k sens. for point sources]

Phase 2: 200 BEACON stations + 3k GRAND antennas [$\approx$ GRAND, IC-Gen2 sens. for diffuse flux]

HERON overview:

K. Kotera, July 16



GRAND + BEACON Collaborations

K. Kotera+1078 | A. Zeolla 1227

GRAND-type:
standalone antennas
15 around each station
autonomous & external triggering
reconstruction



BEACON-type:
phased stations
24 antennas per station
low energy threshold for triggering




HERON: Hybrid Elevated Radio Observatory for Neutrinos

BEACON phased-

Narrow field of
Phase 1: 24 BEACON
Phase 2: 200 BEACON

HERON overview
K. Kotera, July 1



ering)

es]
e flux]

1-2
kms

station

threshold for triggering


GRAND + BEACON Collabor

K. Kotera+1078 | A. Zeolla 1227

PBR: POEMMA-Balloon with Radio

EUSO-SPB design
(UV fluorescence + Cherenkov),
augmented with PUEO-like
radio antennas



PBR overview:
J. Eser, July 22



PBR/JEM-EUSO Collaborations

J. Eser | F. Cafagna | V. Scotti | A. V. Olinto | Z. Plebaniak | M. Bertaina | M. Battisti | L. Wiencke | E. Mayotte | R. Caruso

PBR: POEMMA-Balloon with Radio



EUSO-SPB design

(UV + fluorescence + Cherenkov),
equipped with PUEO-like
radio antennas

PBR overview:
J. Eser, July 22

Anthal
(As)

cosmic Rays $E > \text{PeV}$

EAS

Neutrino
+Radio

Tau Neutrino

PBR/JEM-EUSO Collaborations

J. Eser | F. Cafagna | V. Scotti | A. V. Olinto | Z. Plebaniak | M. Bertaina | M. Battisti | L. Wiencke | E. Mayotte | R. Caruso

What did I think
10 years ago?

What we know / don't know

What we know

- ▶ compatible with isotropy
- ▶ power-law $\propto E^{-2.5}$
- ▶ not coincident with transient sources (*e.g.*, GRBs)
- ▶ not correlated with known sources
- ▶ **flavor composition: compatible with equal proportion of ν_e , ν_μ , ν_τ**
- ▶ also: no prompt atmospheric neutrinos

What we don't know

- ▶ what are the sources?
- ▶ what is the production mechanism?
- ▶ is there a cut-off at 2 PeV?
- ▶ what is the Galactic contribution, if any?
- ▶ what is the precise relation to UHE cosmic rays?
- ▶ is there new physics?
- ▶ **what is the precise flavor composition of the flux?**

...but we have good ideas on all

What we know / don't know

What we know

Still true (2310.15224)

Still true

TXS 0506+056 (+ TDEs?)

NGC 1068

Still true, more precise

Still true

- ▶ compatible with isotropy
- ▶ power-law $\propto E^{-2.5}$
- ▶ not coincident with transient sources (e.g., GRBs)
- ▶ not correlated with known sources
- ▶ **flavor composition: compatible with equal proportion of ν_e , ν_μ , ν_τ**
- ▶ also: no prompt atmospheric neutrinos

What we don't know

- ▶ what are the sources?
- ▶ what is the production mechanism?
- ▶ is there a cut-off at 2 PeV?
- ▶ what is the Galactic contribution, if any?
- ▶ what is the precise relation to UHE cosmic rays?
- ▶ is there new physics?
- ▶ **what is the precise flavor composition of the flux?**

... but we have good ideas on all

What we know / don't know

What we know

Still true (2310.15224)

Still true

TXS 0506+056 (+ TDEs?)

NGC 1068

Still true, more precise

Still true

- ▶ compatible with isotropy
- ▶ power-law $\propto E^{-2.5}$
- ▶ not coincident with transient sources (e.g., GRBs)
- ▶ not correlated with known sources
- ▶ **flavor composition: compatible with equal proportion of ν_e, ν_μ, ν_τ**
- ▶ also: no prompt atmospheric neutrinos

What we don't know

- ▶ what are the sources? ← Unknown, but closer
- ▶ what is the production mechanism? ← Still unknown
- ▶ is there a cut-off at 2 PeV? ← Still unknown
- ▶ what is the Galactic contribution, if any? ← About 10% at ~ TeV
- ▶ what is the precise relation to UHE cosmic rays? ← Unknown, but closer
- ▶ is there new physics? ← Not yet, strong limits
- ▶ **what is the precise flavor composition of the flux?** ← Still unknown

... but we have good ideas on all

From a 2015 talk (about neutrino flavor) at U. Cincinnati:

What we know / don't know

What we know

Still true (2310.15224)

Still true

TXS 0506+056 (+ TDEs?)

NGC 1068

Still true, more precise

Still true

- ▶ compatible with isotropy
- ▶ power-law $\propto E^{-2.5}$
- ▶ not coincident with transient sources (e.g., GRBs)
- ▶ not correlated with known sources
- ▶ **flavor composition: compatible with equal proportion of ν_e , ν_μ , ν_τ**
- ▶ also: no prompt atmospheric neutrinos

What we don't know

- ▶ what are the sources? ← Unknown, but closer
- ▶ what is the production mechanism? ← Still unknown
- ▶ is there a cut-off at 2 PeV? ← Still unknown
- ▶ what is the Galactic contribution, if any? ← About 10% at ~ TeV
- ▶ what is the precise relation to UHE cosmic rays? ← Unknown, but closer
- ▶ is there new physics? ← Not yet, strong limits
- ▶ **what is the precise flavor composition of the flux?** ← Still unknown

~~... but we have good ideas on all~~
... but we have made solid progress on all

How it
started

How it's
going

10–20 years
from now



How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν



How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

First sources
First tests of ν physics

How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

First sources
First tests of ν physics

Mature TeV-PeV ν astronomy
EeV ν discovered
Precision PeV ν physics tests

How it
started

How it's
going

10–20 years
from now

First predictions
of high-energy
cosmic ν

PeV ν
discovered

First sources
First tests of ν physics

How do we get there?

Mature TeV-PeV ν astronomy
EeV ν discovered
Precision PeV ν physics tests


Thank you

(And heartfelt apologies for skipping some contributions)

Backup slides

Neutrinos from the Galaxy


Vitagliano, Tamborra, Raffelt, *RMP* 2019



See also: [Beacom & Candia, JCAP 2004](#)

Neutrinos from the Galaxy

Vitagliano, Tamborra, Raffelt, *RMP* 2019




Search for >10 -TeV
astrophysical ν

- Use *muon tracks*
- Pointing accuracy: $\sim 1^\circ$
- Atm. bg. is mostly ν_μ
- Self-veto screens for atm. muons to cut ν bg.

Neutrinos from the Galaxy

Vitagliano, Tamborra, Raffelt, *RMP* 2019



Search for **TeV**
astrophysical ν

Search for **>10-TeV**
astrophysical ν

- ▶ But GP ν are TeV
- ▶ Use **cascades**
- ▶ Atm. ν_e bg. $10\times$ lower
- ▶ Bg.-to-signal: $10^8:1$
- ▶ *Deep learning retains 20 times more events, $2\times$ better angular res.*

- ▶ Use *muon tracks*
- ▶ Pointing accuracy: $\sim 1^\circ$
- ▶ Atm. bg. is mostly ν_μ
- ▶ Self-veto screens for atm. muons to cut ν bg.


Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:

Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)




Quick aside: how to read a ternary plot

Assumes underlying unitarity –
sum of projections on each axis is 1

How to read it:


Follow the tilt of the tick marks

Always in this order: (f_e, f_μ, f_τ)



Theoretically palatable regions: today

NO, upper θ_{23} octant,
NuFit 5.0




Note:

All plots shown are for normal
neutrino mass ordering (NO);
inverted ordering looks similar

Song, Li, Argüelles, MB, Vincent, JCAP 2021

See also: MB, Beacom, Winter, PRL 2015

Theoretically palatable regions: today




Note:

All plots shown are for normal
neutrino mass ordering (NO);
inverted ordering looks similar

Song, Li, Argüelles, MB, Vincent, JCAP 2021

See also: MB, Beacom, Winter, PRL 2015

Theoretically palatable regions: today




Note:

All plots shown are for normal
neutrino mass ordering (NO);
inverted ordering looks similar

Song, Li, Argüelles, MB, Vincent, JCAP 2021

See also: MB, Beacom, Winter, PRL 2015

Theoretically palatable regions: today




Note:

All plots shown are for normal
neutrino mass ordering (NO);
inverted ordering looks similar

Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2021

See also: **MB**, Beacom, Winter, *PRL* 2015

Theoretically palatable regions: today



Note:

All plots shown are for normal
neutrino mass ordering (NO);
inverted ordering looks similar

Song, Li, Argüelles, **MB**, Vincent, *JCAP* 2021

See also: **MB**, Beacom, Winter, *PRL* 2015

BSM models in connection to KM3-230213A

Lorentz-invariance violation

Superluminal neutrinos (2502.09548, 2502.12070, 2502.18256)

Time delay (14 years!) *vs.* gamma rays from a GRB (2502.13093, 2503.14471)

From the muon surviving enhanced $\mu \rightarrow e + \gamma$ while traveling underground (2502.13201)

Decay of heavy dark matter

Decay of 400-PeV DM (2503.00097, 2503.04464, 2503.14332, 2503.18737, 2504.01447)

Heavy scalar decays into sterile ν that decays into active ν (2503.07776)

Sterile-active neutrino transition

Motivated by observation in KM3NeT, but not IceCube (2502.21299)

Primordial black hole evaporation

Possibly with “memory burden” to lengthen PBH life (2502.19245, 2503.19227, 2503.21740)

Mirror neutrons

UHE $n' \rightarrow n \rightarrow \nu$ reconciles heavy UHECR masses with high cosmogenic ν flux (2503.14419)

Do we live in a simulation? (2504.08461)

Lorentz-invariance violation — from superluminal speeds

A superluminal ν loses energy via pair production, *i.e.*,

$$\nu \rightarrow \nu + e^+ + e^-$$


Cohen & Glashow, *PRL* 2011

Excess over light speed: $\delta = c_\nu - 1$

Decay length: $L_{\text{dec}} = c_\nu / \Gamma \propto E^{-5} \delta^{-3}$

Decay width

Demanding that the travel distance $L < 10 L_{\text{dec}}$ sets upper limits on δ



New limit is ~1000 times stronger than previous one from TXS 0506+056

Lorentz-invariance violation — from a GRB association

Amelino-Camelia *et al.*, 2502.13093

GRB emitted neutrinos & photons simultaneously

Time delay induced by dispersion of neutrinos on spacetime foam:


Neutrino energy

$$\Delta t = D(z) \frac{E}{\Lambda} \approx 14 \text{ years}$$

Cosmological expansion


Energy scale of LIV (10^{14} – 10^{15} GeV)

GRB- ν association: 2.4σ
(p -value of 0.015)



Decay of heavy dark matter ($\text{DM} \rightarrow \nu + \nu$)

Fiorillo, Valera, MB, Winter, *PRD* 2023




Decay of heavy dark matter ($\text{DM} \rightarrow \nu + \nu$)

Galactic DM density profile

NFW (1,3,1.5)


Jho, Park, Shin, 2503.18737



Decay of heavy dark matter — supersymmetric




Multi-component DM: heavy (χ , unstable) & lighter ($\tilde{\chi}_-$, stable)

Jho, Park, Shin, 2503.18737



Sterile-active ν transitions

Brdar & Chattopadhyay, 2502.21299




Sterile-active ν transitions

New neutrino-baryon interactions inside Earth (by gauging $U(1)_B$ symmetry)

Relative strength *vs.* standard weak interaction: $\epsilon_{ss} = G_B/(\sqrt{2}G_F)$

For $-\epsilon_{ss} = 150$, transitions are resonant in KM3NeT,
but not in IceCube




Primordial black hole evaporation

Primordial black holes (PBHs) evaporate through Hawking radiation

“Memory burden” effect:
quantum back-reaction lengthens
the life of the black hole

Most of the contribution is from
intermediate-mass PBHs,
transitioning to memory burden

Galactic + extragalactic contributions,
monochromatic mass spectrum,
PBHs make up all of DM






Mirror neutrons

Can reconcile large cosmogenic ν flux
inspired by KM3-230213A and heavy
UHECR mass composition

But cannot explain lack of IceCube events

Joint fits to Auger UHECR data + neutrino
data from IceCube and KM3NeT



Mirror neutrons

Pierre Auger scenario (2) | $S(z) \propto (1+z)^m$ | $m = 0$ | $z_{\max} = 3$

